

#### CLIA Waived

EZ CLICK® Drug of Abuse Test Cup is a rapid test for the qualitative detection of D-Amphetamine 1000, D-Amphetamine 500, Benzoylecgonine 300, Benzoylecgonine 150, 11-nor-Δ<sup>b</sup>-THC-9-COOH, Oxazepam, Methamphetamine 1000, Methamphetamine 500, Morphine 2000, Methadone, Phencyclidine, Oxycodone, Butablital, Buprenorphine, Morphine 300, 2-Ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine, Methylenedioxymethamphetamine, and Nortriptyline in human urine at a cutoff concentration indicated in the table below.

The test may yield preliminary positive results when prescription drugs are ingested at prescribed doses. It is not intended to distinguish between prescription use and abuse of any drug. There are no uniformly recognized cutoff concentration levels for any drug in urine. The test provides only preliminary test results. A more specific alternative chemical method must be used in order to obtain a confirmed analytical result. GC/MS is the preferred confirmatory method. Clinical consideration and professional judgment should be exercised with any drug of abuse test result, particularly when the preliminary result is positive.

The test may come with adulterant tests [specimen validity tests] of (Specific Gravity (S.G.), Oxidants (OX), pH, Creatinine (CR), Nitrite (NI) and Glutaraldehyde (GL)) for the determination of diluted or adulterated urine specimens.

For in vitro diagnostic use only.

#### WHAT IS EZ CLICK® DRUG OF ABUSE TEST CUP?

EZ CLICK® Drug of Abuse Test Cup is a rapid test for qualitative detection of D-Amphetamine 1000, D-Amphetamine 500, Benzoylecgonine 300, Benzoylecgonine 150, 11-nor-∆-THC-9-COOH, Oxazepam, Methamphetamine 1000, Morphine 2000, Methadone, Oxycodone, Phencyclidine, Butalbital, Buprenorphine, Morphine 300, 2-Ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine, Methylenedioxymethamphetamine, and Northyline in human urine. The EZ CLICK® Drug of Abuse Test Cup yields a positive result when drug and/or its metabolite in urine is at or exceeds its cutoff concentration.

## WHAT IS THE CUT-OFF VALUE AND APPROXIMATE DETECTION TIME?

Drug (Identifier)	Cutoff Level	Minimum Detection Time	Maximum Detection Time
D-Amphetamine (AMP)	1000ng/mL	4-6 hours	2-3 days
D-Amphetamine (AMP 500)	500ng/mL	4-6 hours	2-3 days
Benzoylecgonine (COC)	300ng/mL	2-6 hours	2-3 days
Benzoylecgonine (COC 150)	150ng/mL	2-6 hours	2-3 days
11-nor-Δ <sup>9</sup> -THC-9-COOH (THC)	50ng/mL	1-3 hours	1-7 days
Oxazepam (BZO)	300ng/mL	2-7 hours	1-4 days
Methamphetamine (mAMP)	1000ng/mL	4-6 hours	2-3 days
Methamphetamine (mAMP 500)	500ng/mL	4-6 hours	2-3 days
Morphine (OPI)	2000ng/mL	2-6 hours	1-3 days
Methadone (MTD)	300ng/mL	3-8 hours	1-3 days
Oxycodone (OXY)	100ng/mL	1-3 hours	1-2 days
Phencyclidine (PCP)	25ng/mL	4-6 hours	7-14 days
Butalbital (BARB)	300ng/mL	2-4 hours	1-3 weeks
Buprenorphine (BUP)	10ng/mL	2-6 hours	2-4 days
Morphine (MOR)	300ng/mL	2-6 hours	1-3 days
2-Ethylidene-1,5-dimethyl-3,3- diphenylpyrrolidine (EDDP)	300ng/mL	3-8 hours	1-3 days
Methylenedioxymethamphetamine (MDMA)	500ng/mL	2-7 hours	2-4 days
Nortriptyline (TCA)	1,000ng/mL	8-12 hours	2-7 days

#### PRINCIPLE

The EZ CLICK® Drug of Abuse Test Cup is an immunoassay. During testing, a urine specimen migrates upward on the test strip. A drug-positive urine specimen will not generate a colored line in the specific test line region of the strip, while a drug-negative urine specimen will generate a line in the test line region. A colored line will always appear at the control line region, indicating that proper volume of specimen has been added.

### ADULTERATION SUMMARY AND EXPLANATION OF THE TEST

The adulterant test strip contains chemically treated reagent pads. Observation of the color change on the strip compared to the color chart provides a semi-quantitative screen for oxidants, specific gravity, pH, creatinine, nitrite and glutaraldehyde in human urine which can help to assess the integrity of the urine specimen.

Adulteration is the tampering of a urine specimen with the intention of altering the test results. The use of adulterants in the urine specimen can cause false negative results by either interfering with the test and/or destroying the drugs present in the urine. Dilution may also be used to produce false negative drug test results. To determine certain urinary characteristics and to detect the presence of specific gravity, pH, oxidants, nitrite, glutaraldehyde and creatinine in urine are considered the best ways to test for adulteration or dilution.

- Oxidants (OX): Tests for the presence of oxidizing agents such as bleach and peroxide in the urine.
   Specific Gravity (S.G.): Tests for sample dilution. Normal levels for specific gravity will range from 1.003
- Specific Gravity (S.G.): Tests for sample dilution. Normal levels for specific gravity Will range from 1.003 to 1.030. Specific gravity levels of less than 1.003 or higher than 1.030 may be an indication of adulteration or specimen dilution.
- pH: Tests for the presence of acidic or alkaline adulterants in urine. Normal pH levels should be in the
- range of 4.0 to 9.0. Values below pH 4.0 or above pH 9.0 may indicate the sample has been altered.

  Nitrite (NIT): Tests for commercial adulterants such as Klear and Whizzies. Normal urine specimens
- should contain no trace of nitrite. Positive results for nitrite usually indicate the presence of an adulterant.

   Glutaraldehyde (GLU): Tests for the presence of an aldehyde. Glutaraldehyde is not normally found in a urine specimen. Detection of glutaraldehyde in a specimen is generally an indicator of adulteration.
- Creatinine (CRE): Creatinine is one way to check for dilution and flushing, which are the most common mechanisms used in an attempt to circumvent drug testing. Low creatinine may indicate dilute urine.

#### WARNINGS AND PRECAUTIONS

- For in vitro diagnostic use
- For external use only.
- B For single use Discard after first use
- Do not use the test if the pouch is punctured or not well sealed.
- Do not use after expiration date.
- Keep out of the reach of children.
- . The used test cup and urine specimen should be discarded according to federal, state and local regulations.

#### CONTENT OF THE PACKAGE

Included in package:

- User Instruction
- Test Cup (inside foil pouch)
- Security Seal Label
- Color Chart Card for Adulterants Interpretation (when applicable)

#### Not included in package:

- Watch. Timer or Clock

#### STORAGE AND STABILITY

Store as packaged in the sealed pouch at 39°F - 86°F (4°C - 30°C). The test is stable through the expiration date printed on the sealed pouch. The test cup must remain in the sealed pouch until use. Keep away from direct sunlight, moisture and heat. DO NOT FREEZE. Do not use beyond the expiration date.

#### WHEN TO COLLECT URINE FOR THE TEST?


Urine from any time of day can be use. The minimum detection time varies for different drugs. (Refer to the approximate detection timetable).

#### HOW TO COLLECT URINE?

- When you are ready to begin, remove the test cup from the sealed foil pouch. Peel back and remove
  the label from the test cup to show the drug test strips. Notice the colored tape on each strip correlates
  to the name of the drug you are testing for.
- Remove the cap from the test cup. Fill the test cup with a minimum of 30 mL (see the minimum line mark) fresh urine sample. Do not over-fill (the maximum-line mark).
- When finished, recap the test cup (be sure to tighten it firmly by turning the lid until it clicks) and place the test cup on a flat surface. Be sure NOT to tilt or flip it over.

#### HOW TO DO THE TEST

 After filling the test cup with a fresh urine sample, wait for 5 minutes (start timing immediately after sample is collected) and read the results. DO NOT read results after 5 minutes.





Note: Results after more than 5 minutes may be not accurate and should not be read.

#### READING THE RESULTS

#### Preliminary positive (+)

If a line appears in the C - Control area but NO line appears in the T - Drug Test area, then it indicates a Preliminary Positive result for the corresponding drug.


#### Negative (-)

If a line appears in both the C - Control and T - Drug Test area, then it indicates a Negative result for the corresponding drug regardless of how dark or light the line may appear.

#### Invali

If at 5 minutes, NO line appears in the C - Control area, then the results are Invalid.

Note: Each test strip needs to be look at individually. Each line may vary in color and darkness or the rate at which the line appears. (DO NOT compare lines within the same test strip or between different test extract)



A positive test result does not always mean a person took illegal drugs and a negative test result does not always mean a person did not take illegal drugs. There are a number of factors that influence the reliability of drug tests. Certain drugs of abuse tests are more accurate than others.

**IMPORTANT:** The result you obtained is called preliminary for a reason. The sample must be tested by laboratory in order to determine if a drug of abuse is actually present.

## WHAT IS A FALSE POSITIVE TEST?

The definition of a false positive test would be an instance where the test result from the **EZ CLICK® Drug** of **Abuse Test Cup** is positive, even though the initial target drug is not present in the sample. The most common causes of a false positive test are cross reactants. Certain foods and medicines, diet plan drugs and nutritional supplements may also cause a false positive test result with this product.

#### WHAT IS A FALSE NEGATIVE TEST?

The definition of a false negative test is that the initial target drug is present but is not detect by **EZ CLICK® Drug of Abuse Test Cup.** If the sample is diluted, or if the sample is tainted or contaminated with a substance this could cause false negative results.

#### READING THE ADULTERANT RESULTS

(Please refer to the color chart card)

Semi-quantitative results are obtained by visually comparing the reacted color blocks on the strip to the printed color indicator on the color chart. No instrumentation is required.

#### EST LIMITATIONS

- The EZ CLICK® Drug of Abuse Test Cup provides only a qualitative, preliminary analytical result. A secondary analytical method must be use to obtain a confirmed result. Gas chromatography/mass spectropatry (CCMS) is the preferred confirmation method.
- spectrometry (GC/MS) is the preferred confirmatory method.

  2. There is a possibility that interfering substances in the urine specimen may cause erroneous results.
- Substances, such as bleach and/or alum, in urine specimens may produce erroneous results.
- A positive result does not indicate intoxication, the concentration of drug in the urine, or the route of drug administration.
- A negative result may not necessarily indicate drug-free urine. Negative results can be obtained when drug is present but below the cutoff level of the test.
- Test does not distinguish between drugs of abuse and certain medications.
- 7. A positive test result may be obtained from certain foods or food supplements.

#### ADULTERANT TESTS (SPECIMEN VALIDITY TESTS) LIMITATIONS

- The adulterant tests included with the product are meant to aid in the determination of abnormal specimens but may not cover all the possible adulterants.
- Oxidants: Normal human urine should not contain oxidants. The presence of high level of antioxidants in the specimen, such as ascorbic acid, may result in false negative results for the oxidants pad.
- 3. Specific Gravity: Elevated levels of protein in urine may cause abnormally high specific gravity values.

  A Nitritie: Nitritie is not a normal component of human urine. However, nitrite found in urine may indicate
- Nitrite: Nitrite is not a normal component of human urine. However, nitrite found in urine may indicate urinary tract infections or bacterial infections. Nitrite levels of > 20mg/dL may produce false positive olutaraldehyde results.
- Glutaraldehyde: Glutaraldehyde is normally not found in a urine specimen. However, certain metabolic abnormalities such as ketoacidosis (fasting, uncontrolled diabetes or high-protein diets) may interfere with the test results.
- Creatinine: Creatinine tests for the specimen for dilution and flushing. Normal creatinine levels are between 20 and 350mg/dL. Under rare conditions, certain kidney diseases may show dilute urine.

#### QUALITY CONTROL

If you work in a laboratory, you should perform quality control testing and you should read this section.

A procedural control is included in the test. A color line appearing in the control region (C) is considering an internal procedural control. It confirms sufficient specimen volume, adequate membrane wicking and correct procedural technique.

Control standards are not supply with this kit. However, it is recommended that positive and negative controls be tested as good laboratory practice to confirm the test procedure and to verify proper test performance. Quality control testing should be done with each new lot and each new shipment. It should be done every thirty days to check storage. Please contact our Technical Support at 1-626-443-8480 for controls that work with the test cup.

#### PERFORMANCE CHARACTERISTICS

Eighty clinical urine specimens were analyzed by GC/MS and by the **EZ CLICK® Drug of Abuse Test Cup.** Each test was read by three viewers. Samples were divided by concentration into five categories: drug-free, less than half the cutoff, near cutoff negative, near cutoff positive, and high positive. Results were as follows:

## Accuracy - D-Amphetamine 1000

Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

Viewer R

Viewer B:					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	12	26
Negative	10	10	19	2	0

<sup>%</sup> agreement among positives is 95% % agreement among negatives is 97.5%

r . . . . . . 0

Viewer C:					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

% agreement among negatives is 100%

The average negative agreement is 98.3%.

From the results of the above tables, the total results are shown as below for D-Amphetamine 1000: The average positive agreement is 96.7%.

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

#### Accuracy - D-Amphetamine 500

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

viewei b.					
Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	14	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 100%

From the results of the above tables, the total results are shown as below for D-Amphetamine 500: The average positive agreement is 98.3%.

The average negative agreement is 98.3%

# Accuracy - Benzoylecgonine 300

viewer A:					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	12	26
Negative	10	10	19	2	0

<sup>%</sup> agreement among positives is 95%

## Viewer B:

Orug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
0	0	0	12	26
10	10	20	2	0
	0	trug-Free Cutoff Concentration by GC/MS Analysis  0 0	Less Than Half the Cutoff	Less Than Half the Cutoff Cutoff Concentration by GC/MS Analysis

VICWCI O.					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Benzoylecgonine 300:

The average positive agreement is 95.8%. The average negative agreement is 98.3%

#### Accuracy - Benzoylecgonine 150

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Retween 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	14	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 100% % agreement among negatives is 97.5%

Viower B

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

Viousor C

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Benzoylecgonine 150: The average positive agreement is 98.3%.

The average negative agreement is 98.3%.

# Accuracy - 11-nor-Δ<sup>9</sup>-THC-9-COOH

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	Concentration)	Concentration)	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 100% % agreement among negatives is 97.5%

viewer B:					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	14	26
Negative	10	10	20	0	0

<sup>%</sup> agreement among positives is 100% % agreement among negatives is 100%

VICWCI O.					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	14	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

From the results of the above tables, the total results are shown as below for 11-nor- $\Delta^{g}$ -THC-9-COOH: The average positive agreement is 99.2%.

The average negative agreement is 99.2%.

#### Accuracy - Oxazepam

	WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
ſ	Positive	0	0	0	14	25
ſ	Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	14	25
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

Viewer C:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Retween 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	High Positive (Greater Than 50% above the Cutoff Concentration)
Positive	0	0	0	13	25
Negative	10	10	20	2	0

<sup>%</sup> agreement among positives is 95%

From the results of the above tables, the total results are shown as below for Oxazepam:

The average positive agreement is 96.7%.

The average negative agreement is 100%.

## Accuracy - Methamphetamine 1000

Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	18	21
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

Viewer B

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	18	21
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Retwoon 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%			
Positive	0	0	0	17	21			
Negative	10	10	20	2	0			
% agreeme	% agreement among positives is 95%							

% agreement among negatives is 100.0%

From the results of the above tables, the total results are shown as below for Methamphetamine 1000: The average positive agreement is 96.7%.

The average negative agreement is 100%.

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among positives is 95% % agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 100%

#### Accuracy - Methamphetamine 500

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

#### Viower B

VICWCI D.					
Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Methamphetamine 500: The average positive agreement is 97.5%.

The average negative agreement is 98.3%

#### Accuracy - Morphine 2000

#### Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	15	24
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	15	24
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

VICWCI O.					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	15	24
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Morphine 2000:

The average positive agreement is 97.5%.

#### The average negative agreement is 100%.

#### Accuracy - Methadone

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Retwoon 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

#### Viousor C

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	12	26
Negative	10	10	19	2	0

<sup>%</sup> agreement among positives is 95%

From the results of the above tables, the total results are shown as below for Methadone: The average positive agreement is 96.7%.

The average negative agreement is 97.5%.

# Accuracy - Oxycodone Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	12	26
Negative	10	10	19	2	0

<sup>%</sup> agreement among positives is 95%

## Viewer B

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	2	13	26
Negative	10	10	18	1	0
% agreeme	nt among no	eitivoe ie 97 5%			

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 95%

#### Viewer C

VICWCI O.					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	12	26
Negative	10	10	20	2	0

<sup>%</sup> agreement among positives is 95% % agreement among negatives is 100%

From the results of the above tables, the total results are shown as below for Oxycodone:

The average positive agreement is 95.8%.

#### The average negative agreement is 97.5%.

#### Accuracy - Phencyclidine

	WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Γ	Positive	0	0	2	13	26
Γ	Negative	10	10	18	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 95%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	12	26
Negative	10	10	20	2	0

<sup>%</sup> agreement among positives is 95% % agreement among negatives is 100%

## Viewer C:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Phencyclidine: The average positive agreement is 96.7%.

The average negative agreement is 98.3%.

## Accuracy - Butalbital

#### Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	13	26
Negative	10	10	20	1	0
0/ narnama	nt among no	oitivos is 07 E9/			

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Retween 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	13	26
Negative	10	10	19	1	0
/		aiti			

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Butalbital: The average positive agreement is 97.5%.

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is100 %

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

The average negative agreement is 99.2%.

#### **Accuracy - Buprenorphine**

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

VIEWEI D.	Viewei B.							
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Potycon 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Creater Than 50%			
Positive	0	0	1	13	26			
Negative	10	10	19	1	0			

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

viewer C:					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Buprenorphine:

The average positive agreement is 97.5%.

The average negative agreement is 98.3%.

#### Accuracy - Morphine 300

#### Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

## Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	13	26
Negative	10	10	20	1	0
0/		. '.' ' . 07 50/		•	•

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff	
Positive	0	0	Concentration)	Concentration)	26
Negative	10	10	20	0	0

<sup>%</sup> agreement among positives is 100%

From the results of the above tables, the total results are shown as below for Morphine 300:

The average positive agreement is 98.3%.

#### The average negative agreement is 99.2%

#### Accuracy - 2-Ethylidene-1,5-Dimethyl-3,3-Diphenylpyrrolidine

VIEWEI A.	Wei A.							
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%			
Positive	0	0	1	13	26			
Negative	10	10	19	1	0			

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 100%

viewei C.			Near Cutoff		
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for 2-Ethylidene-1,5-Dimethyl-3,3-Diphenylpyrrolidine:

The average positive agreement is 97.5%.

#### Accuracy - Methylenedioxymethamphetamine Viewer A:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis		Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	14	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 100% % agreement among negatives is 97.5%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

### Viewer C:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	(Potwoon 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Creater Than 50%
Positive	0	0	1	14	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 100%

From the results of the above tables, the total results are shown as below for

#### Accuracy - Nortriptyline

VICWCI 71.					
WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	13	26
Negative	10	10	19	1	0

<sup>%</sup> agreement among positives is 97.5% % agreement among negatives is 97.5%

#### Viewer B:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Near Cutoff Negative (Between 50% below the Cutoff and the Cutoff Concentration)	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	(Greater Than 50%
Positive	0	0	1	14	26
Negative	10	10	19	0	0

<sup>%</sup> agreement among positives is 100% % agreement among negatives is 97.5%

#### Viewer C:

WHPM Result	Drug-Free	Less Than Half the Cutoff Concentration by GC/MS Analysis	Negative (Retween 50%	Near Cutoff Positive (Between the Cutoff and 50% above the Cutoff Concentration)	
Positive	0	0	0	13	26
Negative	10	10	20	1	0

<sup>%</sup> agreement among positives is 97.5%

From the results of the above tables, the total results are shown as below for Nortriptyline: The average positive agreement is 98.3%.

## Precision and Sensitivity - D-Amphetamine 1000

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	2/48
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

# Lot 2

LUI Z		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	3/47
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

# Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	1/49
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 97.5%

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

The average negative agreement is 98.3%.

<sup>%</sup> agreement among negatives is 100%

<sup>%</sup> agreement among negatives is 97.5%

Methylenedioxymethamphetamine: The average positive agreement is 99.2%. The average negative agreement is 98.3%.

<sup>%</sup> agreement among negatives is 100%

The average negative agreement is 98.3%.

# Precision and Sensitivity - D-Amphetamine 500

LOUI		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	3/47
625	50	0/50
750	50	0/50
875	50	0/50
1000	50	0/50

Lot 2		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	2/48
625	50	0/50

50

50

50

0/50

0/50

0/50

ot 3		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	2/48
625	50	0/50
750	50	0/50
875	50	0/50
1000	E0.	0/50

# Precision and Sensitivity - Benzoylecgonine 300 Lot 1

750

875

1000

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

Lot 2		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	2/48
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

# Precision and Sensitivity - Benzoylecgonine 150

Ot 1		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
37.5	50	50/0
75	50	50/0
112.5	50	50/0
150	50	3/47
187.5	50	0/50
225	50	0/50
262.5	50	0/50
300	50	0/50

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
37.5	50	50/0
75	50	50/0
112.5	50	50/0
150	50	3/47
187.5	50	0/50
225	50	0/50
262.5	50	0/50
300	50	0/50

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
37.5	50	50/0
75	50	50/0
112.5	50	50/0
150	50	3/47
187.5	50	0/50
225	50	0/50
262.5	50	0/50
300	50	0/50

# Precision and Sensitivity - 11-nor-Δ<sup>9</sup>-THC-9-COOH

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
12.5	50	50/0
25	50	50/0
37.5	50	50/0
50	50	2/48
62.5	50	0/50
75	50	0/50
87.5	50	0/50
100	50	0/50

_ot 2		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
12.5	50	50/0
25	50	50/0
37.5	50	50/0
50	50	1/49
62.5	50	0/50
75	50	0/50
87.5	50	0/50
100	50	0/50

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
12.5	50	50/0
25	50	50/0
37.5	50	50/0
50	50	2/48
62.5	50	0/50
75	50	0/50
87.5	50	0/50
100	50	0/50

# Precision and Sensitivity - Oxazepam Lot 1

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

Lot 2		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

Lot 3		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	4/46
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

# Precision and Sensitivity - Methamphetamine 1000

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	3/47
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	2/48
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	3/47
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

# Precision and Sensitivity - Methamphetamine 500 Lot 1

LULI		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	2/48
625	50	0/50
750	50	0/50
875	50	0/50
1000	E0	0/50

#### Int 2

.UL Z		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	3/47
625	50	0/50
750	50	0/50
875	50	0/50
1000	50	0/50

#### Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	2/48
625	50	0/50
750	50	0/50
875	50	0/50
1000	50	0/50

# Precision and Sensitivity - Morphine 2000

Lot 1		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
500	50	50/0
1000	50	50/0
1500	50	50/0
2000	50	2/48
2500	50	0/50
3000	50	0/50
3500	50	0/50
4000	50	0/50

# Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
500	50	50/0
1000	50	50/0
1500	50	50/0
2000	50	3/47
2500	50	0/50
3000	50	0/50
3500	50	0/50
4000	50	0/50

#### ot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
500	50	50/0
1000	50	50/0
1500	50	50/0
2000	50	3/47
2500	50	0/50
3000	50	0/50
3500	50	0/50
4000	50	0/50

# Precision and Sensitivity - Methadone

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

## Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

# Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

# Precision and Sensitivity - Oxycodone Lot 1

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
25	50	50/0
50	50	50/0
75	50	50/0
100	50	3/47
125	50	0/50
150	50	0/50
175	50	0/50
200	50	0/50

# Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
25	50	50/0
50	50	50/0
75	50	50/0
100	50	3/47
125	50	0/50
150	50	0/50
175	50	0/50
200	50	0/50

#### Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
25	50	50/0
50	50	50/0
75	50	50/0
100	50	2/48
125	50	0/50
150	50	0/50
175	50	0/50
200	50	0/50

# Precision and Sensitivity - Phencyclidine Lot 1

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
6.3	50	50/0
12.5	50	50/0
18.8	50	50/0
25	50	3/47
31.3	50	0/50
37.5	50	0/50
43.8	50	0/50
50	50	0/50

## Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
6.3	50	50/0
12.5	50	50/0
18.8	50	50/0
25	50	3/47
31.3	50	0/50
37.5	50	0/50
43.8	50	0/50
50	50	0/50

## Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
6.3	50	50/0
12.5	50	50/0
18.8	50	50/0
25	50	3/47
31.3	50	0/50
37.5	50	0/50
43.8	50	0/50
50	50	0/50

#### Precision and Sensitivity - Butalbital Lot 1

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

# Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

#### Lot 3

LUI 3		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

# Precision and Sensitivity - Buprenorphine

_0( )		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
2.5	50	50/0
5	50	50/0
7.5	50	50/0
10	50	3/47
12.5	50	0/50
15	50	0/50
17.5	50	0/50
20	50	0/50

#### Lot 2

JL Z		
Approximate concentration of sample (ng/mL)	Number of determinations	Results Negative/ Positive
0	50	50/0
2.5	50	50/0
5	50	50/0
7.5	50	50/0
10	50	2/48
12.5	50	0/50
15	50	0/50
17.5	50	0/50
20	50	0/50

#### Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
2.5	50	50/0
5	50	50/0
7.5	50	50/0
10	50	3/47
12.5	50	0/50
15	50	0/50
17.5	50	0/50
20	50	0/50

## Precision and Sensitivity - Morphine 300

LULI		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	2/48
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

## Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

#### ot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	2/48
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

#### Precision and Sensitivity - 2-Ethylidene-1,5-Dimethyl-3,3-Diphenylpyrrolidine

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	2/48
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

#### Lot 2

.UL Z		
Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	2/48
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

## Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
75	50	50/0
150	50	50/0
225	50	50/0
300	50	3/47
375	50	0/50
450	50	0/50
525	50	0/50
600	50	0/50

## Precision and Sensitivity - Methylenedioxymethamphetamine

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	3/47
625	50	0/50
750	50	0/50
875	50	0/50
1000	50	0/50

#### Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	2/48
625	50	0/50
750	50	0/50
875	50	0/50
1000	50	0/50

#### Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
125	50	50/0
250	50	50/0
375	50	50/0
500	50	2/48
625	50	0/50
750	50	0/50
875	50	0/50
1000	50	0/50

#### Precision and Sensitivity - Nortriptyline

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	2/48
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

# Lot 2

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	3/47
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

## Lot 3

Approximate Concentration of Sample (ng/mL)	Number of Determinations	Results Negative/Positive
0	50	50/0
250	50	50/0
500	50	50/0
750	50	50/0
1000	50	3/47
1250	50	0/50
1500	50	0/50
1750	50	0/50
2000	50	0/50

## SPECIFICITY AND CROSS REACTIVITY

To test the specificity of the test, the test device was used to test D-Amphetamine 1000, D-Amphetamine 500, Benzoylecgonine 300, Benzoylecgonine 150, 11-nor-0-THC-9-COOH, Oxazepam, Methamphetamine 1000, Methamphetamine 500, Morphine 2000, Methadone, Phencyclidine, Oxycodone, Butalbital, Buprenorphine, Morphine 300, 2-Ethylidene-1,5-Dimethyl-3,3-Diphenylpyrrolidine, Methylenedioxymethamphetamine, Nortriptyline, drug metabolites and other components of the same class that are likely to be present in urine. All the components were added to drug-free normal human urine. The following structurally related compounds produced positive results with the test when tested at levels equal to or greater than the concentrations listed below.

D-Amphetamine 1000	Result
(D-Amphetamine, Cutoff = 1000ng/mL)	Positive at 1000ng/mL
L-Amphetamine	Positive at 100000ng/mL
DL-Amphetamine	Positive at 500ng/mL
(+/-)-3,4-Methylenedioxyamphetamine (MDA)	Positive at 1300ng/mL
Phentermine	Positive at 100000ng/mL
Apomorphine	Positive at 50000ng/mL
β-Phenylethylamine	Positive at 25000ng/mL
Tyramine	Positive at 10000ng/mL
Tryptamine	Positive at 25000ng/mL
D-Methamphetamine	>100000
L-Methamphetamine	>100000
Ephedrine	>100000
3,4-Methylenedioxyethylamphetamine (MDE)	>100000

D-Amphetamine 500	Result
(D-Amphetamine, Cutoff = 500 ng/mL)	Positive at 500ng/mL
L-Amphetamine	Positive at 60000ng/mL
DL-Amphetamine	Positive at 1000ng/mL
Methylenedioxyamphetamine (MDA)	Positive at 600ng/mL
R-(-)-Apomorphine	Positive at 13000ng/mL
β-Phenylethylamine	Positive at 8000ng/mL
Tyramine	Positive at 5000ng/mL
Tryptamine	Positive at 100000ng/mL
Hydroxyamphetamine	Positive at 600ng/mL
D-Pseudoephedrine	Negative at ≥ 10 <sup>5</sup> ng/mL
D-Methamphetamine	Negative at ≥ 10 <sup>5</sup> ng/mL
L-Methamphetamine	Negative at ≥ 10 <sup>5</sup> ng/mL

(±)-Methamphetamine	Negative at ≥ 10 <sup>5</sup> ng/mL
Ephedrine	Negative at ≥10 <sup>5</sup> ng/mL
3,4-Methylenedioxy-N-ethylamphetamine (MDE)	Negative at ≥ 10 <sup>5</sup> ng/mL
3,4-Methylenedioxymethamphetamine (MDMA)	Negative at ≥ 10 <sup>5</sup> ng/mL
Phentermine	Negative at ≥ 10 <sup>5</sup> ng/mL

Benzoylecgonine 300	Result
(Benzoylecgonine, Cutoff = 300ng/mL)	Positive at 300ng/mL
Cocaine Hydrochloride	Positive at 500ng/mL
Cocaethylene	>100000
Ecgonine	>100000

Benzoylecgonine 150	Result
(Benzoylecgonine, Cutoff = 150ng/mL)	Positive at 150ng/mL
Cocaine Hydrochloride	Positive at 3000ng/mL
Norcocaine	Negative at ≥ 10 <sup>5</sup> ng/mL
Cocaethylene	Negative at ≥ 10 <sup>5</sup> ng/mL
Ecgonine	Negative at ≥ 10 <sup>5</sup> ng/mL

11-nor-Δ <sup>9</sup> -THC-9-COOH	Result
(11-nor-Δ9-THC-9-COOH, Cutoff = 50ng/mL)	Positive at 50ng/mL
11-hydroxy-Δ9-Tetrahydrocannabinol	Positive at 15000ng/mL
Δ <sup>8</sup> -Tetrahydrocannabinol	Positive at 8000ng/mL
Δ <sup>9</sup> -Tetrahydrocannabinol	Positive at 7000ng/mL
Cannabinol	>200000
Cannabidiol	>200000

Oxazepam	Result
(Oxazepam, Cutoff = 300ng/mL)	Positive at 300ng/mL
Alprazolam	Positive at 125ng/mL
α-Hydroxyalprazolam	Positive at 2500ng/mL
Bromazepam	Positive at 1565ng/mL
Chlordiazepoxide	Positive at 1560ng/mL
Clobazam	Positive at 65ng/mL
Clonazepam	Positive at 10000ng/mL
Clorazepate Dipotassium	Positive at 195ng/mL
Delorazepam	Positive at 1560ng/mL
Desalkylflurazepam	Positive at 1565ng/mL
Diazepam	Positive at 115ng/mL
Estazolam	Positive at 165ng/mL
Flunitrazepam	Positive at 166ng/mL
Midazolam	Positive at 6500ng/mL
Nitrazepam	Positive at 300ng/mL
Norchlordiazepoxide	Positive at 250ng/mL
Nordiazepam	Positive at 400ng/mL
Temazepam	Positive at 100ng/mL
Triazolam	Positive at 2500 ng/mL
DL-Lorazepam	Negative at ≤ 10 <sup>5</sup> ng/mL
Methamphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL
Morphine	Negative at ≤ 10 <sup>5</sup> ng/mL

Methamphetamine 1000	Result
(D-Methamphetamine, Cutoff = 1000ng/mL)	Positive at 1000ng/mL
(+/-)-3,4-Methylenedioxy-n-ethylamphetamine (MDEA)	Positive at 41600ng/mL
DL-Methamphetamine	Positive at 1000ng/mL
p-Hydroxymethamphetamine	Positive at 27000ng/mL
(+/-)-3,4-Methylenedioxymethamphetamine (MDMA)	Positive at 8000ng/mL
L-Methamphetamine	Positive at 10000ng/mL
Trimethobenzamide	Negative at ≤ 10 <sup>5</sup> ng/mL
Chloroquine	Negative at ≤ 10 <sup>5</sup> ng/mL
Ephedrine	Negative at ≤ 10 <sup>5</sup> ng/mL
Fenfluramine	Negative at ≤ 10 <sup>5</sup> ng/mL
Procaine (Novocaine)	Negative at ≤ 10 <sup>5</sup> ng/mL
Ranitidine (Zantac)	Negative at ≤ 10 <sup>5</sup> ng/mL
D-Amphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL
L-Amphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL
Oxazepam	Negative at ≤ 10 <sup>5</sup> ng/mL
Morphine	Negative at ≤ 10 <sup>5</sup> ng/mL

Methampheta	amine 500	Result
(D-Methamph	etamine, Cutoff = 500 ng/mL)	Positive at 500ng/mL
(±)3,4-Methyle	enedioxy-n-ethylamphetamine (MDEA)	Positive at 20000ng/mL
(±)-Methamph	netamine	Positive at 1000ng/mL
p-Hydroxymet	hamphetamine	Positive at 16000ng/mL

(±)3,4-MDMA	Positive at 2000ng/mL
L-Methamphetamine	Positive at 5000ng/mL
Fenfluramine	Positive at 40000ng/mL
L-Amphetamine	Positive at 60000ng/mL
D-Pseudoephedrine	Negative at ≥ 10 <sup>5</sup> ng/mL
Trimethobenzamide	Negative at ≥ 10 <sup>5</sup> ng/mL
Chloroquine	Negative at ≥ 10 <sup>5</sup> ng/mL
Ephedrine	Negative at ≥ 10 <sup>5</sup> ng/mL
Procaine (Novocaine)	Negative at ≥ 10 <sup>5</sup> ng/mL
Ranitidine (Zantac)	Negative at ≥ 10 <sup>5</sup> ng/mL
D-Amphetamine	Negative at ≥ 10 <sup>5</sup> ng/mL
Oxazepam	Negative at ≥ 10 <sup>5</sup> ng/mL
Morphine	Negative at ≥ 10 <sup>5</sup> ng/mL
(±) 3,4-MDA	Negative at ≥ 10 <sup>5</sup> ng/mL

Morphine 2000	Result
(Morphine, Cutoff = 2000ng/mL)	Positive at 2000ng/mL
Codeine	Positive at 1000ng/mL
Ethylmorphine	Positive at 560ng/mL
Hydrocodone	Positive at 5000ng/mL
Hydromorphone	Positive at 7315ng/mL
Levorphanol	Positive at 16000ng/mL
6-Monoacetylmorphine	Positive at 1000ng/mL
Morphine 3-β-D-Glucuronide	Positive at 1300ng/mL
Thebaine	Negative at ≤ 10 <sup>5</sup> ng/mL
Norcodeine	Negative at ≤ 10 <sup>5</sup> ng/mL
Normorphine	Negative at ≤ 10 <sup>5</sup> ng/mL
Oxycodone	Negative at ≤ 10 <sup>5</sup> ng/mL
Oxymorphone	Negative at ≤ 10 <sup>5</sup> ng/mL
Procaine	Negative at ≤ 10 <sup>5</sup> ng/mL
Oxazepam	Negative at ≤ 10 <sup>5</sup> ng/mL
Methamphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL

Methadone	Result
(Methadone, Cutoff = 300ng/mL)	Positive at 300ng/mL
Levo-α-Acetylmethadol	Positive at 10000ng/mL
Alphamethadol	Negative at ≤ 10 <sup>5</sup> ng/mL
Doxylamine	Negative at ≤ 10 <sup>5</sup> ng/mL
2-Ethylidene-1,5-Dimethyl-3,3-Diphenylpyrrolidine	Negative at ≤ 10 <sup>5</sup> ng/mL
2-Ethyl-5-Methyl-3,3-Diphenylpyrroline	Negative at ≤ 10 <sup>5</sup> ng/mL

Oxycodone	Result
(Oxycodone, Cutoff = 100ng/mL)	Positive at 100ng/mL
Dihydrocodeine	Positive at 50,000ng/mL
Hydrocodone	Positive at 10,000ng/mL
Heroin	Negative at ≤ 10 <sup>5</sup> ng/mL
Morphine 3-β-D-Glucuronide	Negative at ≤ 10 <sup>5</sup> ng/mL
Codeine	Negative at ≤ 10 <sup>5</sup> ng/mL
Hydromorphone	Negative at ≤ 10 <sup>5</sup> ng/mL
Morphine	Negative at ≤ 10 <sup>5</sup> ng/mL
Acetylmorphine	Negative at ≤ 10 <sup>5</sup> ng/mL
Buprenorphine	Negative at ≤ 10 <sup>5</sup> ng/mL
Ethylmorphine	Negative at ≤ 10 <sup>5</sup> ng/mL

Phencyclidine	Result
(Phencyclidine, Cutoff = 25ng/mL)	Positive at 25ng/mL
Phencyclidine Morpholine	Positive at 625ng/mL
4-Hydroxyphencyclidine	Positive at 250ng/mL

Butalbital	Result	
(Butalbital, Cutoff = 300ng/mL)	Positive at 300ng/mL	
Secobarbital	Positive at 300ng/mL	
Amobarbital	Positive at 3000ng/mL	
Alphenal	Positive at 250ng/mL	
Aprobarbital	Positive at 200ng/mL	
Allobarbital	Positive at 500ng/mL	
Butabarbital	Positive at 1000ng/mL	
Butethal	Positive at 500ng/mL	
Cyclopentobarbital	Positive at 300ng/mL	
Pentobarbital	Positive at 1300ng/mL	
Phenobarbital	Positive at 1900ng/mL	

Result

Positive at 10ng/mL

Buprenorphine

(Buprenorphine, Cutoff = 10ng/mL)

Buprenorphine-3-D-Glucuronide	Positive at 15ng/mL
Norbuprenorphine	Positive at 40ng/mL
Norbuprenorphine-3-D-Glucuronide	Positive at 500ng/mL
Morphine	Negative at ≤ 10 <sup>5</sup> ng/mL
Oxymorphone	Negative at ≤ 10 <sup>5</sup> ng/mL

Morphine 300	Result		
(Morphine, Cutoff = 300ng/mL)	Positive at 300ng/mL		
6-Acetylmorphine	Positive at 750ng/mL		
Codeine	Positive at 300ng/mL		
Ethylmorphine	Positive at 200ng/mL		
Heroin	Positive at 700ng/mL		
Hydromorphone	Positive at 4000ng/mL		
Hydrocodone	Positive at 2000ng/mL	Positive at 2000ng/mL	
Levorphanol	Positive at 12000ng/mL		
Thebaine	Positive at 90000ng/mL		
Methyprylon	Positive at 4000ng/mL		
Morphine-3-β-Glucuronide	Positive at 450ng/mL		
Oxycodone	Negative at ≤ 10 <sup>5</sup> ng/mL		
Procaine	Negative at ≤ 10 <sup>5</sup> ng/mL		

2-Ethylidene-1,5-Dimethyl-3,3- Diphenylpyrrolidine	Result
(2-Ethylidene-1,5-Dimethyl-3,3-Diphenylpyrrolidine, Cutoff = 300ng/mL)	Positive at 300ng/mL
2-Ethyl-5-Methyl-3,3-Diphenylpyrroline (EMDP)	Negative at ≤ 10 <sup>5</sup> ng/mL
Disopyramide	Negative at ≤ 10 <sup>5</sup> ng/mL
Methadone	Negative at ≤ 10 <sup>5</sup> ng/mL
Levo-α-Acetylmethadol Hydrochloride (LAAM)	Negative at ≤ 10 <sup>5</sup> ng/mL
Alphamethadol	Negative at ≤ 10 <sup>5</sup> ng/mL
Doxylamine	Negative at ≤ 10 <sup>5</sup> ng/mL

Methylenedioxymethamphetamine	Result
(Methylenedioxymethamphetamine, Cutoff = 500ng/mL)	Positive at 500ng/mL
3,4-Methylenedioxyamphetamine HCI (MDA)	Positive at 8000ng/mL
3,4-Methylenedioxyethylamphetamine (MDEA)	Positive at 1000ng/mL
(-)-Ψ-Ephedrine	Positive at 40000ng/mL
D-Methamphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL
D-Amphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL
L-Amphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL
L-Methamphetamine	Negative at ≤ 10 <sup>5</sup> ng/mL

Nortriptyline	Result
(Nortriptyline, Cutoff = 1000ng/mL)	Positive at 1000ng/mL
Amitriptyline	Positive at 1500ng/mL
Clomipramine	Positive at 15000ng/mL
Desipramine	Positive at 1000ng/mL
Doxepin	Positive at 2000ng/mL
Imipramine	Positive at 600ng/mL
Nordoxepin	Positive at 1000ng/mL
Promazine	Positive at 24000ng/mL
Trimipramine	Positive at 4000ng/mL
Cyclobenzaprine Hydrochloride	Positive at 1500ng/mL
Maprotiline	Negative at ≤ 10 <sup>5</sup> ng/mL
Promethazine	Negative at ≤ 10 <sup>5</sup> ng/mL
Norclomipramine	Negative at ≤ 10 <sup>5</sup> ng/mL

#### EFFECT OF URINARY SPECIFIC GRAVITY

Urine samples of normal, high, and low specific gravity ranges (1.000-1.035) were spiked with drugs at 25% below and 25% above cutoff levels respectively. **EZ CLICK® Drug of Abuse Test Cup** was tested using twelve drug-free urine and spiked urine samples. The results demonstrate that varying ranges of urinary specific gravity do not affect the test results.

#### EFFECT OF URINARY PH

The pH of an aliquot of negative urine pool was adjusted to pH ranges of 4.0 - 9.0 and spiked with drugs at 25% below and 25% above cut-off levels. The spiked, pH-adjusted urine was tested with *EZ CLICK® Drug of Abuse Test Cup*. The results demonstrate that varying ranges of pH do not interfere with the performance of the test.

Non Cross-Reacting Compounds - D-Amphetamine 1000

4-Acetamidophenol	L-Ephedrine	Oxycodone		
Acetophenetidin	(-)-Ψ-Ephedrine	Oxymetazoline		
N-Acetylprocainamide	Erythromycin	Papaverine		
Acetylsalicylic Acid	β-Estradiol	Penicillin-G		
Aminopyrine	Estrone-3-Sulfate	Pentazocine		
Amitriptyline	Ethyl-p-Aminobenzoate	Pentobarbital		
Amobarbital	Fenfluramine	Perphenazine		
Amoxicillin	Fenoprofen	Phencyclidine		

Ampicillin	Furosemide	Phenelzine
Ascorbic Acid	Gentisic Acid	Phenobarbital
Aspartame	Hemoglobin	Phenytoin
Atropine	Hydralazine	L-Phenylephrine
Benzilic Acid	Hydrochlorothiazide	Phenylpropanolamine
Benzoic Acid	Hydrocodone	Prednisolone
Benzoylecgonine	Hydrocortisone	Prednisone
Bilirubin	o-Hydroxyhippuric Acid	Procaine
Brompheniramine	3-Hydroxytyramine	Promazine
Caffeine	Ibuprofen	Promethazine
Cannabidiol	Imipramine	DL-Propranolol
Cannabinol	(-)-Isoproterenol	D-Propoxyphene
Chloral Hydrate	Isoxsuprine	Quinidine
Chloramphenicol	Ketamine	Quinine
Chlordiazepoxide	Ketoprofen	Ranitidine
Chlorothiazide	Labetalol	Salicylic Acid
(±)-Chlorpheniramine	Levorphanol	Secobarbital
Chlorpromazine	Loperamide	Sulfamethazine
Chloroquine	Maprotiline	Sulindac
Cholesterol	Meperidine	Temazepam
Clomipramine	Meprobamate	Tetracycline
Clonidine	Methadone	Tetrahydrocortisone
Cocaine Hydrochloride	Methylphenidate	Tetrahydrozoline
Codeine	Morphine 3-β-D-Glucuronide	Δ9-THC-COOH
Cortisone	Nalidixic Acid	Thebaine
(-)-Cotinine	Naloxone	Thiamine
Creatinine	Naltrexone	Thioridazine
Deoxycorticosterone	Naproxen	DL-Thyroxine
Dextromethorphan	Niacinamide	Tolbutamide
Diazepam	Nifedipine	Triamterene
Diclofenac	Norcodeine	Trifluoperazine
Diflunisal	Norethindrone	Trimethoprim
Digoxin	D-Norpropoxyphene	Trimipramine
Diphenhydramine	Noscapine	Tryptamine
Doxylamine	DL-Octopamine	DL-Tyrosine
Ecgonine Hydrochloride	Oxalic Acid	Uric Acid
Ecgonine Methyl Ester	Oxazepam	Verapamil
(IR,2S)-(-)-Ephedrine	Oxolinic Acid	Zomepirac

Non	Cross-Reacting	Compounds	- D-Ami	ohetamine 500	

Acetaminophen (4- Acetamidophenol)	Erythromycin	Oxalic Acid
Acetophenetidin	β-Estradiol	Oxazepam
N-Acetylprocainamide	Fenoprofen	Oxolinic Acid
Acetylsalicylic Acid	Furosemide	Oxymetazoline
Albumin	Gentisic Acid	Papaverine
Aminopyrine	Hemoglobin	Penicillin-G
D-Amphetamine	Hydralazine	Pentobarbital
Amoxicillin	Hydrochlorothiazide	Perphenazine
Ampicillin	Hydrocodone	Phenelzine
Apomorphine	Hydrocortisone	Phencyclidine
L-Ascorbic Acid	o-Hydroxyhippuric Acid	Prednisone
Aspartame	3-Hydroxytyramine	Procaine
Atropine	Ibuprofen	DL-Propranolol
Benzilic Acid	D,L-Isoproterenol	D-Propoxyphene
Benzoic Acid	Isoxsuprine	D-Pseudoephedrine
Benzoylecgonine	Ketamine	Quinine
Bilirubin	Ketoprofen	Ranitidine
Cannabidiol	Labetalol	Salicylic Acid
Chloral Hydrate	Loperamide	Secobarbital
Chloramphenicol	Maprotiline	Serotonin (5-Hydroxytyramine)
Chlorothiazide	Meperidine	Sulfamethazine
Chlorpromazine	Meprobamate	Sulindac
Chloroquine	Methadone	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cholesterol	Methamphetamine	Tetrahydrozoline
Clonidine	Methoxyphenamine	Thiamine
Codeine	Morphinie-3-β-D-Glucuronide	Thioridazine
Cortisone	Nalidixic acid	Triamterene
(-)-Cotinine	Naloxone	DL-Tyrosine
Creatinine	Naltrexone	Trifluoperazine
Deoxycorticosterone	Naproxen	Trimethoprim
Dextromethorphan	Niacinamide	DL-Tryptophan
Diclofenac	Nifedipine	Tyramine
Diflunisal	Norcodeine	Uric Acid

Digoxin	Norethindrone	Verapamil
Diphenhydramine	D-Norpropoxyphene	Zomepirac
Ecgonine Methyl Ester	Noscapine	
EMDD	DL-Octopamine	

# Non Cross-Reacting Compounds - Benzoylecgonine 300

Non Cross-Reacting Compounds		
Acetaminophen	Estrone-3-Sulfate	Oxymetazoline
Acetophenetidin	Ethyl-p-Aminobenzoate	Papaverine
N-Acetylprocainamide	Fenoprofen	Penicillin-G
Acetylsalicylic Acid	Furosemide	Pentobarbital
Aminopyrine	Gentisic Acid	Perphenazine
Amitriptyline	Hemoglobin	Phencyclidine
Amobarbital	Hydralazine	Phenelzine
Amoxicillin	Hydrochlorothiazide	Phenobarbital
Ampicillin	Hydrocodone	Phentermine
L-Ascorbic Acid	Hydrocortisone	L-Phenylephrine
DL-Amphetamine Sulfate	o-Hydroxyhippuric Acid	β-Phenylethylamine
Apomorphine	p-Hydroxymethamphetamine	Phenylpropanolamine
Aspartame	3-Hydroxytyramine	Prednisolone
Atropine	Ibuprofen	Prednisone
Benzilic Acid	Imipramine	Procaine
Benzoic Acid	Iproniazid	Promazine
Benzphetamine	(±)-Isoproterenol	Promethazine
(±)-Brompheniramine	Isoxsuprine	DL-Propranolol
Caffeine	Ketamine	D-Propoxyphene
Cannabidiol	Ketoprofen	D-Pseudoephedrine
Cannabinol	Labetalol	Quinidine
Chloral Hydrate	Levorphanol	Quinine
Chloramphenicol	Loperamide	Ranitidine
Chlordiazepoxide	Maprotiline	Salicylic Acid
Chlorothiazide	Meperidine	Secobarbital
(±)-Chlorpheniramine	Meprobamate	Serotonin
Chlorpromazine	Methadone	Sulfamethazine
Chloroquine	Methoxyphenamine	Sulindac
,	(±)-3,4-	
Cholesterol	Methylenedioxyamphetamine	Temazepam
Clomipramine	(±)-3,4- Methylenedioxymethamphetamine Hydrochloride	Tetracycline
Clonidine	Morphine-3-β-D-Glucuronide	Tetrahydrocortisone 3-(β-D- Glucuronide)
Codeine	Morphine Sulfate	Tetrahydrozoline
Cortisone	Nalidixic acid	Thebaine
(-)-Cotinine	Naloxone	Thiamine
Creatinine	Naltrexone	Thioridazine
Deoxycorticosterone	Naproxen	DL-Tyrosine
Dextromethorphan	Niacinamide	Tolbutamide
Diazepam	Nifedipine	Triamterene
Diclofenac	Norcodeine	Trifluoperazine
Diflunisal	Norethindrone	Trimethoprim
Digoxin	D-Norpropoxyphene	Trimipramine
Diphenhydramine	Noscapine	Tryptamine
Doxylamine	DL-Octopamine	DL-Tryptophan
Ecgonine Methyl Ester	Oxalic Acid	Tyramine
(-)-Ψ-Ephedrine	Oxazepam	Uric Acid
Erythromycin	Oxolinic Acid	Verapamil
β-Estradiol	Oxycodone	Zomepirac
F ========	,	

# Non Cross-Reacting Compounds - Benzoylecgonine 150

Acetaminophen (4- Acetamidophenol)	Erythromycin	Oxalic Acid
Acetophenetidin	β-Estradiol	Oxazepam
N-Acetylprocainamide	Fenoprofen	Oxolinic Acid
Acetylsalicylic Acid	Furosemide	Oxymetazoline
Albumin	Gentisic Acid	Papaverine
Aminopyrine	Hemoglobin	Penicillin-G
D-Amphetamine	Hydralazine	Pentobarbital
Amoxicillin	Hydrochlorothiazide	Perphenazine
Ampicillin	Hydrocodone	Phenelzine
Apomorphine	Hydrocortisone	Phencyclidine
L-Ascorbic Acid	o-Hydroxyhippuric Acid	Prednisone
Aspartame	3-Hydroxytyramine	Procaine
Atropine	Ibuprofen	DL-Propranolol
Benzilic Acid	D,L-Isoproterenol	D-Propoxyphene
Benzoic Acid	Isoxsuprine	D-Pseudoephedrine
Benzoylecgonine	Ketamine	Quinine

Bilirubin	Ketoprofen	Ranitidine
Cannabidiol	Labetalol	Salicylic Acid
Chloral Hydrate	Loperamide	Secobarbital
Chloramphenicol	Maprotiline	Serotonin (5-Hydroxytyramine)
Chlorothiazide	Meperidine	Sulfamethazine
Chlorpromazine	Meprobamate	Sulindac
Chloroquine	Methadone	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cholesterol	Methamphetamine	Tetrahydrozoline
Clonidine	Methoxyphenamine	Thiamine
Codeine	Morphinie-3-β-D-Glucuronide	Thioridazine
Cortisone	Nalidixic acid	Triamterene
(-)-Cotinine	Naloxone	DL-Tyrosine
Creatinine	Naltrexone	Trifluoperazine
Deoxycorticosterone	Naproxen	Trimethoprim
Dextromethorphan	Niacinamide	DL-Tryptophan
Diclofenac	Nifedipine	Tyramine
Diflunisal	Norcodeine	Uric Acid
Digoxin	Norethindrone	Verapamil
Diphenhydramine	D-Norpropoxyphene	Zomepirac
Ecgonine Methyl Ester	Noscapine	
EMDP	DL-Octopamine	

#### Non Cross-Reacting Compounds - 11-nor-Δ9-THC-9-COOH

	inds - 11-nor-Δ³-THC-9-COOH	In .
4-Acetamidophenol	β-Estradiol	Papaverine
Acetophenetidin	Estrone-3-Sulfate	Penicillin-G
N-Acetylprocainamide	Ethyl-p-Aminobenzoate	Pentazocine
Acetylsalicylic Acid	Fenoprofen	Pentobarbital
Aminopyrine	Furosemide	Perphenazine
Amitriptyline	Gentisic Acid	Phencyclidine
Amobarbital	Hemoglobin	Phenelzine
Amoxicillin	Hydralazine	Phenobarbital
Ampicillin	Hydrochlorothiazide	Phentermine
Ascorbic Acid	Hydrocodone	L-Phenylephrine
DL-Amphetamine	Hydrocortisone	β-Phenethylamine
L-Amphetamine	o-Hydroxyhippuric Acid	β-Phenylethylamine
Apomorphine	3-Hydroxytyramine	Phenylpropanolamine
Aspartame	Ibuprofen	Prednisolone
Atropine	Imipramine	Prednisone
Benzilic Acid	Iproniazid	Procaine
Benzoic Acid	(-)-Isoproterenol	Promazine
Benzoylecgonine	Isoxsuprine	Promethazine
Benzphetamine	Ketamine	DL-Propranolol
Bilirubin	Labetalol	D-Propoxyphene
Brompheniramine	Levorphanol	D-Pseudoephedrine
Caffeine	Loperamide	Quinidine
Chloral Hydrate	Maprotiline	Quinine
Chloramphenicol	Meprobamate	Ranitidine
Chlordiazepoxide	Methadone	Salicylic Acid
Chlorothiazide	Methoxyphenamine	Secobarbital
	(+)-3.4-	
(±)-Chlorpheniramine	Methylenedioxyamphetamine	Serotonin (5-Hydroxytyramine)
Chlorpromazine	(+)-3,4- Methylenedioxymethamphetamine	Sulfamethazine
Chloroquine	Methylphenidate	Sulindac
Cholesterol	Methyprylon	Temazepam
Clomipramine	Morphine 3-β-D-Glucuronide	Tetracycline
Clonidine	Nalorphine	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cocaine Hydrochloride	Naloxone	Tetrahydrozoline
Codeine	Nalidixic Acid	Thebaine
Cortisone	Naltrexone	Thiamine
(-)-Cotinine	Naproxen	Thioridazine
Creatinine	Niacinamide	DL-Thyroxine
Deoxycorticosterone	Nifedipine	Tolbutamide
Dextromethorphan	Norcodeine	Triamterene
Diazepam	Norethindrone	Trifluoperazine
Diclofenac	D-Norpropoxyphene	Trimethoprim
Diflunisal	Noscapine	Trimipramine
Digoxin	DL-Octopamine	Tryptamine
Diphenhydramine	Oxalic Acid	DL-Tryptophan
Doxylamine	Oxazepam	Tyramine
Ecgonine Hydrochloride	Oxolinic Acid	DL-Tyrosine
Ecgonine Methyl Ester	Oxycodone	Uric Acid
(-)-Ψ-Ephedrine	Oxymetazoline	Verapamil
Erythromycin	p-Hydroxymethamphetamine	Zomepirac

Non Cross-Reacting	Compounds - Oxazepam
--------------------	----------------------

Non Cross-Reacting Compo	ounds - Oxazepam	
4-Acetamidophenol	Diphenhydramine	DL-Octopamine
Acetophenetidin	Doxylamine	Oxalic Acid
N-Acetylprocainamide	Ecgonine Hydrochloride	Oxolinic Acid
Acetylsalicylic Acid	Ecgonine Methyl Ester	Pentobarbital
Aminopyrine	(-)-Ψ-Ephedrine	Perphenazine
Amitriptyline	Fenoprofen	Phencyclidine
Amobarbital	Furosemide	Phenelzine
Amoxicillin	Gentisic Acid	Phenobarbital
Ampicillin	Hemoglobin	Phentermine
L-Ascorbic Acid	Hydrocortisone	L-Phenylephrine
DL-Amphetamine	o-Hydroxyhippuric Acid	β-Phenylethylamine
Apomorphine	p-Hydroxymethamphetamine	Phenylpropanolamine
Aspartame	3-Hydroxytyramine	Prednisone
Atropine	Ibuprofen	DL-Propranolol
Benzilic Acid	Imipramine	D-Pseudoephedrine
Benzoic Acid	(±)-Isoproterenol	Ranitidine
Benzoylecgonine	Isoxsuprine	Salicylic Acid
Benzphetamine	Ketamine	Secobarbital
Bilirubin	Ketoprofen	Serotonin (5-Hydroxytyramine)
(±)-Chlorpheniramine	Labetalol	Sertraline
Caffeine	Loperamide	Sulfamethazine
Cannabidiol	Maprotiline	Sulindac
Chloral Hydrate	Meperidine	Tetrahydrocortisone 3-(β-D- Glucuronide)
Chloramphenicol	Meprobamate	Tetrahydrozoline
Chlorothiazide	Methadone	Thiamine
Chlorpromazine	Methoxyphenamine	Thioridazine
Chloroquine	(+)-3,4- Methylenedioxyamphetamine	DL-Tyrosine
Cholesterol	(+)-3,4- Methylenedioxymethamphetamine	Tolbutamide
Clomipramine	Nalidixic Acid	Triamterene
Clonidine	Nalorphine	Trifluoperazine
Cocaine Hydrochloride	Naloxone	Trimethoprim
Cortisone	Naltrexone	Tryptamine
(-)-Cotinine	Naproxen	DL-Tryptophan
Creatinine	Niacinamide	Tyramine
Dextromethorphan	Nifedipine	Uric Acid
Diclofenac	Norethindrone	Verapamil
Diflunisal	D-Norpropoxyphene	Zomepirac
Dioxin	Noscapine	
Iproniazid	Quinine	

# Non Cross-Reacting Compounds - Methamphetamine 1000

Acetaminophen	Gentisic Acid	Oxycodone
Acetophenetidin	Glucuronide	Oxymetazoline
N-Acetylprocainamide	Glutethimide	Papaverine
Acetylsalicylate	Guaifenesin	Penicillin-G
Aminopyrine	Hippuric Acid	Pentazocine
Amitriptyline	Hydralazine	Pentobarbital
Amobarbital	Hydrochlorothiazide	Perphenazine
Amoxicillin	Hydrocodone	Phencyclidine
Ampicillin	Hydrocortisone	Phenelzine
Apomorphine	o-Hydroxyhippuric Acid	Phenobarbital
Aspartame	3-Hydroxytyramine	Prednisolone
Atropine	Ibuprofen	Phenylpropanolamine
Benzilic Acid	Imipramine	Prednisone
Benzoic Acid	(-)-Isoproterenol	Procaine
Benzoylecgonine	Isoxsuprine	Promazine
Butabarbital	Ketamine	Promethazine
Cannabidiol	Ketoprofen	DL-Propranolol
Chloral Hydrate	Labetalol	D-Propoxyphene
Chloramphenicol	Levorphanol	D-Pseudoephedrine
Chlordiazepoxide	Loperamide	Quinidine
Chlorothiazide	Loxapine Succinate	Quinine
Chlorpromazine	Maprotiline	Ranitidine
Cholesterol	Meperidine	Salicylic Acid
Clomipramine	Meprobamate	Secobarbital
Clonidine	Methadone	Serotonin (5-Hydroxytyramine)
Cocaine Hydrochloride	Methaqualone	Sulfamethazine
Codeine	Methylphenidate	Sulindac
Cortisone	Methyprylon	Temazepam
(-)-Cotinine	Morphine-3-β-D-Glucuronide	Tetracycline
Creatinine	Nalidixic Acid	Tetrahydrocortisone 3-(β-D-

		Glucuronide)
Deoxycorticosterone	Nalorphine	Tetrahydrozoline
Dextromethorphan	Naloxone	Thebaine
Diazepam	Naltrexone	Thiamine
Diclofenac	Naproxen	Thioridazine
Diflunisal	Niacinamide	Tolbutamide
Digoxin	Nifedipine	Triamterene
Diphenhydramine	Norcodeine	Trifluoperazine
Doxylamine	Norethindrone	Trimethoprim
Ecgonine Hydrochloride	Noroxymorphone	Trimipramine
Ecgonine Methyl Ester	D-Norpropoxyphene	DL-Tryptophan
Erythromycin	Noscapine	Tyramine
β-Estradiol	Nylidrin	DL-Tyrosine
Estrone-3-Sulfate	DL-Octopamine	Uric Acid
Ethyl-p-Aminobenzoate	Oxalic Acid	Verapamil
Fenoprofen	Oxazepam	Zomepirac
Furosemide	Oxolinic Acid	

Acetaminophen (4- Acetamidophenol)	Erythromycin	Oxalic Acid
Acetophenetidin	β-Estradiol	Oxazepam
N-Acetylprocainamide	Fenoprofen	Oxolinic Acid
Acetylsalicylic Acid	Furosemide	Oxymetazoline
Albumin	Gentisic Acid	Papaverine
Aminopyrine	Hemoglobin	Penicillin-G
D-Amphetamine	Hydralazine	Pentobarbital
Amoxicillin	Hydrochlorothiazide	Perphenazine
Ampicillin	Hydrocodone	Phenelzine
Apomorphine	Hydrocortisone	Phencyclidine
L-Ascorbic Acid	o-Hydroxyhippuric Acid	Prednisone
Aspartame	3-Hydroxytyramine	Procaine
Atropine	Ibuprofen	DL-Propranolol
Benzilic Acid	D,L-Isoproterenol	D-Propoxyphene
Benzoic Acid	Isoxsuprine	D-Pseudoephedrine
Benzoylecgonine	Ketamine	Quinine
Bilirubin	Ketoprofen	Ranitidine
Cannabidiol	Labetalol	Salicylic Acid
Chloral Hydrate	Loperamide	Secobarbital
Chloramphenicol	Maprotiline	Serotonin (5-Hydroxytyramine)
Chlorothiazide	Meperidine	Sulfamethazine
Chlorpromazine	Meprobamate	Sulindac
Chloroquine	Methadone	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cholesterol	Methamphetamine	Tetrahydrozoline
Clonidine	Methoxyphenamine	Thiamine
Codeine	Morphinie-3-β-D-Glucuronide	Thioridazine
Cortisone	Nalidixic acid	Triamterene
(-)-Cotinine	Naloxone	DL-Tyrosine
Creatinine	Naltrexone	Trifluoperazine
Deoxycorticosterone	Naproxen	Trimethoprim
Dextromethorphan	Niacinamide	DL-Tryptophan
Diclofenac	Nifedipine	Tyramine
Diflunisal	Norcodeine	Uric Acid
Digoxin	Norethindrone	Verapamil
Diphenhydramine	D-Norpropoxyphene	Zomepirac
Ecgonine Methyl Ester	Noscapine	
EMDP	DL-Octopamine	

## Non Cross-Reacting Compounds - Morphine 2000

4-Acetamidophenol	Ecgonine Methyl Ester	Oxolinic Acid
Acetophenetidin	(-)-Ψ-Ephedrine	Oxymetazoline
N-Acetylprocainamide	Erythromycin	Papaverine
Acetylsalicylic Acid	β-Estradiol	Penicillin-G
Aminopyrine	Estrone-3-Sulfate	Pentazocine
Amitriptyline	Ethyl-p-Aminobenzoate	Pentobarbital
Amobarbital	Fenoprofen	Perphenazine
Amoxicillin	Furosemide	Phencyclidine
Ampicillin	Gentisic Acid	Phenelzine
Ascorbic Acid	Hemoglobin	Phenobarbital
DL-Amphetamine	Hydralazine	Phentermine
Apomorphine	Hydrochlorothiazide	L-Phenylephrine
Aspartame	Hydrocortisone	β-Phenylethylamine
Atropine	o-Hydroxyhippuric Acid	Phenylpropanolamine
Benzilic Acid	n-Hydroxymethamphetamine	Prednisone

Benzoic Acid	3-Hydroxytyramine	DL-Propranolol
Benzoylecgonine	Ibuprofen	D-Propoxyphene
Benzphetamine	Imipramine	D-Pseudoephedrine
(±)-Bilirubin	Iproniazid	Quinidine
Brompheniramine	Isoproterenol	Quinine
Caffeine	Isoxsuprine	Ranitidine
Cannabidiol	Ketamine	Salicylic Acid
Chloral Hydrate	Ketoprofen	Secobarbital
Chloramphenicol	Labetalol	Serotonin (5-Hydroxytyramine)
Chlordiazepoxide	Loperamide	Sulfamethazine
Chlorothiazide	Maprotiline	Sulindac
(±)-Chlorpheniramine	Meperidine	Temazepam
Chlorpromazine	Meprobamate	Tetracycline
Chloroquine	Methadone	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cholesterol	Methoxyphenamine	Tetrahydrozoline
Clomipramine	(+)-3,4- Methylenedioxyamphetamine	Thiamine
Clonidine	(+)-3,4- Methylenedioxymethamphetamine	Thioridazine
Cocaine Hydrochloride	Nalidixic Acid	DL-Tyrosine
Cortisone	Nalorphine	Tolbutamide
(-)-Cotinine	Naloxone	Triamterene
Creatinine	Naltrexone	Trifluoperazine
Deoxycorticosterone	Naproxen	Trimethoprim
Dextromethorphan	Niacinamide	Trimipramine
Diazepam	Nifedipine	Tryptamine
Diclofenac	Norethindrone	DL-Tryptophan
Diflunisal	D-Norpropoxyphene	Tyramine
Digoxin	Noscapine	Uric Acid
Diphenhydramine	DL-Octopamine	Verapamil
Doxylamine	Oxalic Acid	Zomepirac
Ecgonine Hydrochloride	Oxazepam	

#### Non Cross-Reacting Compounds - Methadone

Non Cross-Reacting Compo	unds - Methadone	
Acetaminophen	Erythromycin	Oxycodone
Acetophenetidin	β-Estradiol	Oxymetazoline
N-Acetylprocainamide	Estrone-3-Sulfate	Papaverine
Acetylsalicylic Acid	Ethyl-p-Aminobenzoate	Penicillin-G
Aminopyrine	Fenoprofen	Pentazocine Hydrochloride
Amitriptyline	Furosemide	Pentobarbital
Amobarbital	Gentisic Acid	Perphenazine
Amoxicillin	Hemoglobin	Phencyclidine
Ampicillin	Hydralazine	Phenelzine
L-Ascorbic Acid	Hydrochlorothiazide	Phenobarbital
DL-Amphetamine Sulfate	Hydrocodone	Phentermine
Apomorphine	Hydrocortisone	L-Phenylephrine
Aspartame	o-Hydroxyhippuric Acid	β-Phenylethylamine
Atropine	p-Hydroxyamphetamine	Phenylpropanolamine
Benzilic Acid	p-Hydroxymethamphetamine	Prednisolone
Benzoic Acid	3-Hydroxytyramine	Prednisone
Benzoylecgonine	Ibuprofen	Procaine
Benzphetamine	Imipramine	Promazine
Bilirubin	Iproniazid	Promethazine
Caffeine	(±)-Isoproterenol	DL-Propranolol
Cannabidiol	Isoxsuprine	D-Propoxyphene
Cannabinol	Ketamine	D-Pseudoephedrine
Chloral Hydrate	Ketoprofen	Quinacrine
Chloramphenicol	Labetalol	Quinidine
Chlorothiazide	Levorphanol	Quinine
Chlorpromazine	Loperamide	Ranitidine
Chloroquine	Maprotiline	Salicylic Acid
Cholesterol	Meperidine	Secobarbital
Clomipramine	Meprobamate	Serotonin
Clonidine	Methamphetamine	Sulfamethazine
Cocaethylene	Methoxyphenamine	Sulindac
Temazepam	(±)-3,4- Methylenedioxyamphetamine Hydrochloride	Tetracycline
Cocaine Hydrochloride	(±)-3,4- Methylenedioxymethamphetamine Hydrochloride	Tetrahydrocortisone 3-(β-D-Glucuronide)
Codeine	Morphine-3-β-D-Glucuronide	Tetrahydrozoline
Cortisone	Morphine Sulfate	Thebaine
(-)-Cotinine	Nalidixic Acid	Thiamine
Creatinine	Naloxone	Thioridazine

Deoxycorticosterone	Naltrexone	DL-Tyrosine
Dextromethorphan	Naproxen	Tolbutamide
Diazepam	Niacinamide	Triamterene
Diclofenac	Nifedipine	Trifluoperazine
Diflunisal	Norcodeine	Trimethoprim
Digoxin	Norethindrone	Trimipramine
Diphenhydramine	D-Norpropoxyphene	Tryptamine
Ecgonine Hydrochloride	Noscapine	DL-Tryptophan
Ecgonine Methyl Ester	DL-Octopamine	Tyramine
(-)-Ψ-Ephedrine	Oxalic Acid	Uric Acid
(1R,2S)-(-)-Ephedrine	Oxazepam	Verapamil
L-Epinephrine	Oxolinic Acid	Zomepirac

Non Cross-Reacting Compounds - Ovycodone

Non Cross-Reacting Comp		
Acetophenetidin	Ethyl-p-Aminobenzoate	Papaverine
Acetylsalicylic Acid	β-Estradiol	Penicillin-G
Aminopyrine	Estrone-3-Sulfate	Perphenazine
Amoxicillin	Erythromycin	Phenelzine
Ampicillin	Fenoprofen	L-Phenylephrine
Apomorphine	Furosemide	β-Phenylethylamine
Aspartame	Gentisic Acid	Phenylpropanolamine
Atropine	Hemoglobin	Prednisone
Benzilic Acid	Hydralazine	Loperamide
Benzoic Acid	Hydrochlorothiazide	Quinine
Benzphetamine	Hydrocortisone	Quinidine
Bilirubin	o-Hydroxyhippuric Acid	Ranitidine
Deoxycorticosterone	3-Hydroxytyramine	Salicylic Acid
Caffeine	Labetalol	Serotonin
Chloral Hydrate	DL-Isoproterenol	Sulfamethazine
Chloramphenicol	Meprobamate	Sulindac
Chlorothiazide	Methoxyphenamine	Tetracycline
DL-Chlorpheniramine	Nalidixic Acid	Tetrahydrocortisone
Chlorpromazine	Naloxone	Morphine-3-β-D-Glucuronide
Chloroquine	Naltrexone	Tetrahydrozoline
Cholesterol	Naproxen	Thiamine
Clonidine	Niacinamide	Thioridazine
L-Cotinine	Nifedipine	DL-Tyrosine
Cortisone	Isoxsuprine	Tolbutamide
Creatinine	DL-Propranolol	Triamterene
D-Pseudoephedrine	Ketoprofen	Trifluoperazine
Dextromethorphan	Norethindrone	Trimethoprim
Diclofenac	D-Norpropoxyphene	Tyramine
Diflunisal	Noscapine	DL-Tryptophan
Digoxin	DL-Octopamine	Urine Acid
Diphenhydramine	Oxalic Acid	Verapamil
L-Ephedrine	Oxolinic Acid	Zomepirac
Ecgonine Methyl Ester	Oxymetazoline	

Acetaminophen	(-)-Ψ-Ephedrine	Oxycodone
Acetophenetidin	Erythromycin	Oxymetazoline
N-Acetylprocainamide	β-Estradiol	Papaverine
Acetylsalicylic Acid	Estrone-3-Sulfate	Penicillin-G
Aminopyrine	Ethyl-p-Aminobenzoate	Pentazocine Hydrochloride
Amitriptyline	Fenoprofen	Pentobarbital
Amobarbital	Furosemide	Perphenazine
Amoxicillin	Gentisic Acid	Phenelzine
Ampicillin	Hemoglobin	Phenobarbital
Ascorbic Acid	Hydralazine	Phentermine
DL-Amphetamine	Hydrochlorothiazide	L-Phenylephrine
Apomorphine Acid	Hydrocodone	β-Phenylethylamine
Aspartame	Hydrocortisone	Phenylpropanolamine
Atropine	o-Hydroxyhippuric	Prednisolone
Benzilic Acid	p-Hydroxymethamphetamine	Prednisone
Benzoic Acid	3-Hydroxytyramine	Procaine
Benzoylecgonine	Ibuprofen	Promazine
Benzphetamine	Imipramine	Promethazine
Bilirubin	Iproniazid	DL-Propranolol
Brompheniramine	(±)-Isoproterenol	D-Propoxyphene
Caffeine	Isoxsuprine	D-Pseudoephedrine
Cannabidiol	Ketamine	Quinidine
Cannabinol	Ketoprofen	Quinine
Chloral Hydrate	Labetalol	Ranitidine
Chloramphenicol	Loperamide	Salicylic Acid

Chlordiazepoxide	Maprotiline	Secobarbital
Chlorothiazide	Meperidine	Serotonin (5-Hydroxytyramine)
(±)-Chlorpheniramine	Meprobamate	Sulfamethazine
Chlorpromazine	Methadone	Sulindac
Chloroquine	Methoxyphenamine	Temazepam
Cholesterol	(+)-3,4- Methylenedioxyamphetamine	Tetracycline
Clomipramine	(+)-3,4- Methylenedioxymethamphetamine	Tetrahydrocortisone 3-(β-D- Glucuronide)
Clonidine	Morphine-3-β-D-Glucuronide	Tetrahydrozoline
Cocaine Hydrochloride	Morphine Sulfate	Thiamine
Codeine	Nalidixic Acid	Thioridazine
Cortisone	Naloxone	DL-Tyrosine
(-)-Cotinine	Naltrexone	Tolbutamide
Creatinine	Naproxen	Triamterene
Deoxycorticosterone	Niacinamide	Trifluoperazine
Dextromethorphan	Nifedipine	Trimethoprim
Diazepam	Norcodeine	Trimipramine
Diclofenac	Norethindrone	Tryptamine
Diflunisal	D-Norpropoxyphene	DL-Tryptophan
Digoxin	Noscapine	Tyramine
Diphenhydramine	DL-Octopamine	Uric Acid
Doxylamine	Oxalic Acid	Verapamil
Ecgonine Hydrochloride	Oxazepam	Zomepirac
Ecgonine Methyl Ester	Oxolinic Acid	

Non Cross-Reacting Compounds - Butalbital

Non Cross-Reacting Compounds	- Butaibitai	
Acetaminophen	Erythromycin	Nortriptyline
Acetophenetidin	β-Estradiol	o-Hydroxyhippuric Acid
Acetylsalicylic Acid	Estrone-3-Sulfate	DL-Octopamine
Aminopyrine	Ethyl-p-Aminobenzoate	Oxalic Acid
Amitriptyline	Fenoprofen	Oxazepam
Amoxicillin	Furosemide	Oxolinic Acid
Amphetamine	Gentisic Acid	Oxycodone
Ampicillin	Hemoglobin	Oxymetazoline
Apomorphine	Hydralazine	Papaverine
Ascorbic Acid	Hydrochlorothiazide	Penicillin-G
Aspartame	Hydrocodone	Pentazocine
Atropine	Hydrocortisone	Perphenazine
Benzilic Acid	p-Hydroxyamphetamine	Phencyclidine
Benzoic Acid	p-Hydroxymethamphetamine	Phenelzine
Benzoylecgonine	3-Hydroxytyramine	β-Phenethylamine
Bilirubin	Ibuprofen	Phenylpropanolamine
Brompheniramine	Imipramine	Prednisolone
Buprenorphine	(-)-Isoproterenol	Prednisone
Caffeine	Isoxsuprine	Procaine
Cannabidiol	Ketamine	Promazine
Cannabinol	Ketoprofen	Promethazine
Chloral Hydrate	Labetalol	DL-Propranolol
Chloramphenicol	Levorphanol	D-Propoxyphene
Chlorothiazide	Loperamide	Quinidine
(±)-Chlorpheniramine	L-Phenylephrine	Quinine
Chlorpromazine	Maprotiline	Ranitidine
Chloroquine	Meperidine	Salicylic Acid
Cholesterol	Meprobamate	Serotonin
Clomipramine	Morphine	Sulfamethazine
Clonidine	Morphine-3-β-D-Glucuronide	Sulindac
	Methadone	Temazepam
Cocaine Hydrochloride Codeine	Methamphetamine	Tetracycline
Cortisone	(±)-3,4- Methylenedioxyamphetamine Hydrochloride	Tetrahydrozoline
(-)-Cotinine	Methylenedioxymethamphetamine	Thebaine
Creatinine	Morphine Sulfate	Thiamine
Deoxycorticosterone	N-Acetylprocainamide	Thioridazine
Dextromethorphan	Nalidixic Acid	Triamterene
Diazepam	Naloxone	Trifluoperazine
Diclofenac	Naltrexone	Trimethoprim
Diflunisal	Naproxen	Trimipramine
Digoxin	Niacinamide	Tryptamine
Diphenhydramine	Nifedipine	DL-Tyrosine
Doxylamine Doxylamine	Norcodeine	Uric Acid
,	Norcodeine Norethindrone	Verapamil
Ecgonine Hydrochloride		'
Ecgonine Methyl Ester	D-Norpropoxyphene	Zomepirac
(IR,2S)-(-)-Ephedrine	Noscapine	

Ī
---

Non Cross-Reacting Compounds	- Buprenorphine	
4-Acetamidophenol	Erythromycin	Oxolinic Acid
Acetophenetidin	β-Estradiol	Oxycodone
N-Acetylprocainamide	Estrone-3-Sulfate	Oxymetazoline
Acetylsalicylic Acid	Ethyl-p-Aminobenzoate	Papaverine
Aminopyrine	Fenoprofen	Penicillin-G
Amobarbital	Furosemide	Pentazocine Hydrochloride
Amoxicillin	Gentisic Acid	Pentobarbital
Ampicillin	Hemoglobin	Perphenazine
L-Ascorbic Acid	Hydralazine	Phencyclidine
Amphetamine	Hydrochlorothiazide	Phenelzine
Apomorphine	Hydrocodone	Phenobarbital
Aspartame	Hydrocortisone	Phentermine
Atropine	o-Hydroxyhippuric Acid	β-Phenylethylamine
Benzilic Acid	p-Hydroxyamphetamine	Trans-2-Phenylcyclopropylamine Hydrochloride
Benzoic Acid	p-Hydroxymethamphetamine	L-Phenylephrine
Benzoylecgonine	3-Hydroxytyramine	Phenylpropanolamine
Benzphetamine	Ibuprofen	Prednisolone
Bilirubin	Iprazid	Prednisone
(±)-Brompheniramine	(±)-Isoproterenol	Procaine
Butalbital	Isoxsuprine	DL-Propranolol
Caffeine	Ketamine	D-Propoxyphene
Cannabidiol	Ketoprofen	D-Pseudoephedrine
Cannabinol	Labetalol	Quinacrine
Chloral Hydrate	Loperamide	Quinidine
Chloramphenicol	3,4-Methylenedioxy-N- Ethylamphetamine	Quinine
Chlorothiazide	Meperidine	Ranitidine
(±)-Chlorpheniramine	Meprobamate	Salicylic Acid
Chlorpromazine	Methadone	Secobarbital
Chloroquine	L-Methamphetamine	Serotonin
Cholesterol	Methoxyphenamine	Sulfamethazine
Clonidine	(±)-3,4- Methylenedioxyamphetamine Hydrochloride	Sulindac
Cocaethylene	Methylenedioxymethamphetamine	Tetracycline
Cocaine Hydrochloride	Morphine	Tetrahydrocortisone 3-(β-D- Glucuronide)
Codeine	Morphine-3-β-D-Glucuronide	Tetrahydrozoline
Cortisone	Morphine Sulfate	Thiamine
(-)-Cotinine	Nalidixic Acid	Thioridazine
Creatinine	Naloxone	DL-Tyrosine
Deoxycorticosterone	Naltrexone	Tolbutamide
Dextromethorphan	Naproxen	Triamterene
Diclofenac	Niacinamide	Trifluoperazine
Diflunisal	Nifedipine	Trimethoprim
Digoxin	Norcodeine	Tryptamine
Diphenhydramine	Norethindrone	DL-Tryptophan
Doxylamine	D-Norpropoxyphene	Tyramine
Ecgonine Hydrochloride	11-nor-Δ <sup>9</sup> -THC-9-COOH	Uric Acid
Ecgonine Methyl Ester	Nortriptyline	Verapamil
		Zomepirac
Ephedrine	Noscapine	Zomepirac
Ephedrine L-Epinephrine 2-Ethylidene-1,5-Dimethyl-3,3-	Noscapine Oxalic Acid	Zomepirac

Non Cross-Reacting Compounds - Morphine 300

Acebutolol	(-)-Ψ-Ephedrine	Oxymetazoline
Acetylpromazine-d6	Erythromycin	p-Hydroxymethamphetamine
4-Acetamidophenol	β-Estradiol	Papaverine
Acetophenetidin	Estrone-3-Sulfate	Penicillin-G
N-Acetylprocainamide	Ethyl-p-Aminobenzoate	Pentazocine
Acetylsalicylic Acid	2-Ethylidene-1,5-Dimethyl-3,3- Diphenylpyrrolidine	Pentobarbital
Aminopyrine	Fenoprofen	Perphenazine
Amitriptyline	Furosemide	Phencyclidine
Amobarbital	Gentisic Acid	Phenelzine
Amoxicillin	Hemoglobin	Phenobarbital
Ampicillin	Hydralazine	Phentermine
Ascorbic Acid	Hydrochlorothiazide	L-Phenylephrine
Amphetamine	Hydrocortisone	β-Phenethylamine
L-Amphetamine	o-Hydroxyhippuric Acid	β-Phenylethylamine
Apomorphine	3-Hydroxytyramine	Phenylpropanolamine
Aspartame	Ibuprofen	Prednisolone

Atropine	Imipramine	Prednisone
Benzilic Acid	Iprazid	Promazine
Benzoic Acid	(-)-Isoproterenol	Promethazine
Benzoylecgonine	Isoxsuprine	DL-Propranolol
Benzphetamine	Ketamine	D-Propoxyphene
Bilirubin	Ketoprofen	D-Pseudoephedrine
Brompheniramine	Labetalol	Quinidine
Buprenorphine	Loperamide	Quinine
Butalbital	Maprotiline	Ranitidine
Caffeine	Meprobamate	Salicylic Acid
Chloral Hydrate	Methadone	Secobarbital
Chloramphenicol	Methamphetamine	Serotonin (5-Hydroxytyramine)
Chlordiazepoxide	Methoxyphenamine	Sulfamethazine
Chlorothiazide	(+)-3,4- Methylenedioxyamphetamine	Sulindac
(±)-Chlorpheniramine	Methylenedioxymethamphetamine	Temazepam
Chlorpromazine	Methylphenidate	Tetracycline
Chloroquine	Nalorphine	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cholesterol	Naloxone	Tetrahydrozoline
Clomipramine	Nalidixic Acid	Thiamine
Clonidine	Naltrexone	Thioridazine
Cocaine Hydrochloride	Naproxen	DL-Thyroxine
Cortisone	Niacinamide	Tolbutamide
(-)-Cotinine	Nifedipine	Triamterene
Creatinine	Norcodeine	Trifluoperazine
Deoxycorticosterone	Norethindrone	Trimethoprim
Dextromethorphan	D-Norpropoxyphene	Trimipramine
Diazepam	11-nor-Δ9-THC-9-COOH	Tryptamine
Diclofenac	Nortriptyline	DL-Tryptophan
Diflunisal	Noscapine	Tyramine
Digoxin	DL-Octopamine	DL-Tyrosine
Diphenhydramine	Oxalic Acid	Uric Acid
Doxylamine	Oxazepam	Verapamil
Ecgonine Hydrochloride	Oxycodone	Zomepirac
Ecgonine Methyl Ester	Oxolinic Acid	

	unds - 2-Ethylidene-1,5-Dimethyl-3,3	
Acetaminophen	Ecgonine Hydrochloride	o-Hydroxyhippuric Acid
Acetophenetidin	Ecgonine Methyl Ester	Oxalic Acid
Acetylsalicylic Acid	(IR,2S)-(-)-Ephedrine	Oxazepam
Amobarbital	Erythromycin	Oxolinic Acid
Aminopyrine	β-Estradiol	Oxycodone
Amitriptyline	Estrone-3-Sulfate	Oxymetazoline
Amoxicillin	Ethyl p-Aminobenzoate	Papaverine
DL-Amphetamine Sulfate	Fenoprofen	Penicillin-G
Ampicillin	Furosemide	Pentazocine
Apomorphine	Gentisic Acid	Pentobarbital
Ascorbic Acid	Hemoglobin	Perphenazine
Aspartame	Hydralazine	Phencyclidine
Atropine	Hydrochlorothiazide	Phenelzine
Benzilic Acid	Hydrocodone	Phenobarbital
Benzoic Acid	Hydrocortisone	Phentermine
Benzoylecgonine	p-Hydroxyamphetamine	β-Phenylethylamine
Bilirubin	p-Hydroxymethamphetamine	Phenylpropanolamine
Brompheniramine	3-Hydroxytyramine	Prednisolone
Caffeine	Ibuprofen	Prednisone
Cannabidiol	Imipramine	Procaine
Cannabinol	(-)-Isoproterenol	Promazine
Chloral Hydrate	Isoxsuprine	Promethazine
Chloramphenicol	Ketamine	Quinidine
Chlorothiazide	Ketoprofen	Quinine
(±)-Chlorpheniramine	Labetalol	Ranitidine
Chlorpromazine	Levorphanol	Salicylic Acid
Chloroquine	Loperamide	Secobarbital
Cholesterol	L-Phenylephrine	Serotonin
Clomipramine	Maprotiline	Sulfamethazine
Clonidine	Meperidine	Sulindac
Cocaine Hydrochloride	Meprobamate	Temazepam
Codeine	Methamphetamine	Tetracycline
(-)-Cotinine	Methoxyphenamine	Tetrahydrocortisone 3-(β-D- Glucuronide)
Cortisone	(±)-3,4- Methylenedioxyamphetamine Hydrochloride	Tetrahydrozoline
Creatinine	(±)-3,4-	Thebaine

	Methylenedioxymethamphetamine Hydrochloride	
Deoxycorticosterone	Morphine Sulfate	Thiamine
Dextromethorphan	Morphine-3-β-D-Glucuronide	Thioridazine
Diazepam	N-Acetylprocainamide	Triamterene
Diclofenac	Nalidixic Acid	Trifluoperazine
Diflunisal	Naloxone	Trimethoprim
Digoxin	Naltrexone	Trimipramine
Diphenhydramine	Naproxen	Tryptamine
D-Norpropoxyphene	Niacinamide	DL-Tryptophan
D-Propoxyphene	Nifedipine	Tyramine
DL-Tyrosine	Norcodeine	Uric Acid
DL-Octopamine	Norethindrone	Verapamil
DI -Propranolol	Noscanine	Zomenirac

D-Propoxyphene	Nifedipine	Tyramine
DL-Tyrosine	Norcodeine	Uric Acid
DL-Octopamine	Norethindrone	Verapamil
DL-Propranolol	Noscapine	Zomepirac
Non Cross-Reacting Compound	ls - Methylenedioxymethamphetan	nine
4-Acetamidophenol	L-Epinephrine	Pentobarbital
Acetophenetidin	Erythromycin	Perphenazine
N-Acetylprocainamide	β-Estradiol	Phencyclidine
Acetylsalicylic Acid	Estrone-3-Sulfate	Phenelzine
Aminopyrine	Ethyl-p-Aminobenzoate	Phenobarbital
Amitriptyline	Fenoprofen	Phentermine
Amobarbital	Furosemide	Trans-2-Phenylcyclopropylamine Hydrochloride
Amoxicillin	Gentisic Acid	L-Phenylephrine
Ampicillin	Hemoglobin	β-Phenylethylamine
L-Ascorbic Acid	Hydralazine	Phenylpropanolamine
Apomorphine	Hydrochlorothiazide	Prednisolone
Aspartame	Hydrocodone	Prednisone
Atropine	Hydrocortisone	Procaine
Benzilic Acid	o-Hydroxyhippuric Acid	Promazine
Benzoic Acid	3-Hydroxytyramine	Promethazine
Benzoylecgonine	Ibuprofen	DL-Propranolol
Bilirubin	Imipramine	D-Propoxyphene
(±)-Brompheniramine	Iproniazid	D-Pseudoephedrine
Buspirone	(±)-Isoproterenol	Quinacrine
Caffeine	Isoxsuprine	Quinidine
Cannabidiol	Ketamine	Quinine
Cannabinol	Ketoprofen	Ranitidine
Chloral Hydrate	Labetalol	Salicylic Acid
Chloramphenicol	Levorphanol	Secobarbital
Chlordiazepoxide	Loperamide	Serotonin (5-Hydroxytyramine)
Chlorothiazide	Maprotiline	Sulfamethazine
(±)-Chlorpheniramine	Meperidine	Sulindac
Chlorpromazine	Meprobamate	Sustiva
Chloroquine Methylphenidate	Methadone	Temazepam
Cholesterol	Morphine-3-β-D-Glucuronide	Tetracycline
Clomipramine	Morphine Sulfate	Tetrahydrocortisone 3-(β-D- Glucuronide)
Clonidine	Nalidixic Acid	Tetrahydrozoline
Cocaethylene	Naloxone	Thebaine
Cocaine Hydrochloride	Naltrexone	Theophylline
Codeine	Naproxen	Thiamine
Cortisone	Niacinamide	Thioridazine
(-)-Cotinine	Nifedipine	Tolbutamide
Creatinine	Nimesulide	Trazodone
Deoxycorticosterone	Norcodeine	Triamterene
Dextromethorphan	Norethindrone	DL-Tyrosine
Diclofenac	D-Norpropoxyphene	Trifluoperazine
Diazepam	Noscapine	Trimethoprim
Diflunisal	DL-Octopamine	Trimipramine
Digoxin	Oxalic Acid	Tryptamine
Dicyclomine	Oxazepam	DL-Tryptophan
Diphenhydramine	Oxazepani Oxolinic Acid	Tyramine
5,5-Diphenylhydantoin	Oxycodone	Uric Acid
Doxylamine	Oxymetazoline	Verapamil
•	<u> </u>	
Ecgonine Hydrochloride Ecgonine Methyl Ester	Papaverine Penicillin-G	Zomepirac
		+
(IR,2S)-(-)-Ephedrine	Pentazocine Hydrochloride	1

Non Cross-Reacting Compounds - Nortriptyline						
4-Acetamidophenol	Erythromycin	Oxycodone				
Acetophenetidin	β-Estradiol	Oxymetazoline				
N-Acetylprocainamide	Estrone-3-Sulfate	Papaverine				
Acetylsalicylic Acid	Ethyl-p-Aminobenzoate	Penicillin-G				
Aminopyrine	Fenoprofen	Pentazocine Hydrochloride				

Amobarbital	Furosemide	Pentobarbital
Amoxicillin	Gentisic Acid	Perphenazine
Ampicillin	Hemoglobin	Phencyclidine
L-Ascorbic Acid	Hydralazine	Phenelzine
DL-Amphetamine Sulfate	Hydrochlorothiazide	Phenobarbital
Apomorphine	Hydrocodone	Phentermine
Aspartame	Hydrocortisone	β-Phenylethylamine
Atropine	o-Hydroxyhippuric Acid	Trans-2-Phenylcyclopropylamine Hydrochloride
Benzilic Acid	p-Hydroxyamphetamine	L-Phenylephrine
Benzoic Acid	p-Hydroxymethamphetamine	Phenylpropanolamine
Benzoylecgonine	3-Hydroxytyramine	Prednisolone
Benzphetamine	Ibuprofen	Prednisone
Bilirubin	Iproniazid	Procaine
(±)-Brompheniramine	(±)-Isoproterenol	DL-Propranolol
Caffeine	Isoxsuprine	D-Propoxyphene
Cannabidiol	Ketamine	D-Pseudoephedrine
Cannabinol	Ketoprofen	Quinacrine
Chloral Hydrate	Labetalol	Quinidine
Chloramphenicol	Loperamide	Quinine
Chlorothiazide	MDE	Ranitidine
(±)-Chlorpheniramine	Meperidine	Salicylic Acid
Chlorpromazine	Meprobamate	Secobarbital
Chloroquine	Methadone	Serotonin
Cholesterol	L-Methamphetamine	Sulfamethazine
Clonidine	Methoxyphenamine	Sulindac
Cocaethylene	(±)-3,4- Methylenedioxyamphetamine Hydrochloride	Tetracycline
Cocaine Hydrochloride	(+)-3,4- Methylenedioxymethamphetamine Hydrochloride	Tetrahydrocortisone 3-(β-D- Glucuronide)
Codeine	Morphine-3-β-D-Glucuronide	Tetrahydrozoline
Cortisone	Morphine Sulfate	Thiamine
(-)-Cotinine	Nalidixic Acid	Thioridazine
Creatinine	Naloxone	DL-Tyrosine
Deoxycorticosterone	Naltrexone	Tolbutamide
Dextromethorphan	Naproxen	Triamterene
Diclofenac	Niacinamide	Trifluoperazine
Diflunisal	Nifedipine	Trimethoprim
Digoxin	Norcodeine	Tryptamine
Diphenhydramine	Norethindrone	DL-Tryptophan
Doxylamine	D-Norpropoxyphene	Tyramine
Ecgonine Hydrochloride	Noscapine	Uric Acid
Ecgonine Methyl Ester	Oxalic Acid	Verapamil
Ephedrine	Oxazepam	Zomepirac
L-Epinephrine	Oxolinic Acid	,
		l

LAY USER

A lay user study was performed at three intended user sites with 140 laypersons for each drug device. They had diverse educational and professional backgrounds and ranged in age from 21 to >50. Unit samples were prepared at the following concentrations; negative, +/-75%, +/-59%, +/-25% of the cutoff by spiking drug(s) into drug free-pooled urine specimens. The concentrations of the samples were confirmed by GC/MS. Each sample was aliquoted into individual containers and blind-labeled. Each participant was provided with the package insert, 1 blind labeled sample and a device. The results are summarized below.

Ni	Number of	Number of D-Amphetamine 1000	Lay Perso	The Percentage	
% of Cutoff	Samples	Concentration by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	246	0	20	100%
-50% Cutoff	20	492	0	20	100%
-25% Cutoff	20	738	2	18	90%
+25%Cutoff	20	1268	19	1	95%
+50%Cutoff	20	1521	20	0	100%
+75%Cutoff	20	1775	20	0	100%

% of Cutoff Number of Samples	Number of	D-Amphetamine 500 Concentration by GC/MS (ng/mL)	Lay Perso	The Percentage	
			No. of Positive	No. of Negative	Agreement (%)
-100% Cutoff	20	0	0	20	100%
-75% Cutoff	20	130	0	20	100%
-50% Cutoff	20	245	0	20	100%
-25% Cutoff	20	374	0	20	100%
+25% Cutoff	20	649	19	1	95%
+50% Cutoff	20	768	20	0	100%
+75% Cutoff	20	857	20	0	100%
+100% Cutoff	20	1019	20	0	100%

Number of	Benzoylecgonine 300	Lay Pers	The Percentage		
% of Cutoff	Samples	Concentration by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	71	0	20	100%
-50% Cutoff	20	142.5	0	20	100%
-25% Cutoff	20	213.8	2	18	90%
+25%Cutoff	20	379	20	0	100%
+50%Cutoff	20	454.5	20	0	100%
+75%Cutoff	20	530	20	0	100%

	Number of	Benzoylecgonine 150 Concentration by GC/MS (ng/mL)	Lay Perso	The Percentage	
% of Cutoff	Samples		No. of Positive	No. of Negative	Agreement (%)
-100% Cutoff	20	0	0	20	100%
-75% Cutoff	20	37	0	20	100%
-50% Cutoff	20	73	0	20	100%
-25% Cutoff	20	109	1	19	95%
+25% Cutoff	20	178	20	0	100%
+50% Cutoff	20	223	20	0	100%
+75% Cutoff	20	258	20	0	100%
+100% Cutoff	20	301	20	0	100%

% of Cutoff Number of Samples	Number of	11-nor-Δ <sup>9</sup> -THC-9-COOH Concentration by GC/MS (ng/mL)	Lay Person Results		The Percentage
			No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	12	0	20	100%
-50% Cutoff	20	24.5	0	20	100%
-25% Cutoff	20	36.8	1	19	95%
+25%Cutoff	20	64	19	1	95%
+50%Cutoff	20	77	20	0	100%
+75%Cutoff	20	90	20	0	100%

	Number of Oxazepan	Oxazepam Concentration by	Lay Person Results		The Percentage
% of Cutoff	Samples	GC/MS (ng/mL)	No. of Positive	No. of Positive	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	76	0	20	100%
-50% Cutoff	20	145	0	20	100%
-25% Cutoff	20	222	2	18	90%
+25%Cutoff	20	384	20	0	100%
+50%Cutoff	20	468	20	0	100%
+75%Cutoff	20	542	20	0	100%

N	Number of	Methamphetamine 1000	Lay Perso	n Results	The Percentage	
% of Cutoff	Samples	Concentration by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)	
-100%Cutoff	20	0	0	20	100%	
-75%Cutoff	20	245	0	20	100%	
-50% Cutoff	20	488	0	20	100%	
-25% Cutoff	20	729	0	20	100%	
+25%Cutoff	20	1212	19	1	95%	
+50%Cutoff	20	1441	20	0	100%	
+75%Cutoff	20	1666	20	0	100%	

	Number of	Methamphetamine 500	Lay Perso	The Percentage	
% of Cutoff	Samples	Concentration by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100% Cutoff	20	0	0	20	100%
-75% Cutoff	20	129	0	20	100%
-50% Cutoff	20	259	0	20	100%
-25% Cutoff	20	393	1	19	95%
+25% Cutoff	20	669	19	1	95%
+50% Cutoff	20	730	20	0	100%
+75% Cutoff	20	930	20	0	100%
+100% Cutoff	20	1028	20	0	100%

Number of	Number of Samples Morphine 2000 Concentration by GC/MS (ng/mL)	Lay Perso	The Percentage	
		No. of Positive	No. of Negative	Agreement (%)
20	0	0	20	100%
20	527	0	20	100%
20	1053	0	20	100%
20	1573	0	20	100%
20	2652	19	1	95%
20	3254	20	0	100%
20	3711	20	0	100%
	20 20 20 20 20 20 20 20	Samples         by GCMS (ng/mL)           20         0           20         527           20         1053           20         1573           20         2652           20         3254	Number of Samples         Morphine 2000 Concentration by GCMS (ng/mL)         No. of Positive           20         0         0           20         527         0           20         1053         0           20         1573         0           20         2662         19           20         3254         20	Samples         by GC/MS (ng/mL)         No. of Positive Positive         No. of Negative           20         0         0         20           20         527         0         20           20         1053         0         20           20         1573         0         20           20         2652         19         1           20         3254         20         0

	Number of	Methadone Concentration by	Lay Person Results		The Percentage
% of Cutoff	Samples	GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	74	0	20	100%
-50% Cutoff	20	148	0	20	100%
-25% Cutoff	20	222	1	19	95%
+25%Cutoff	20	378	19	1	95%
+50%Cutoff	20	452	20	0	100%
+75%Cutoff	20	530	20	0	100%

	Number of	Oxycodone Concentration by	Lay Perso	The Percentage		
% of Cutoff	Samples			No. of Negative	Agreement (%)	
-100%Cutoff	20	0	0	20	100%	
-75%Cutoff	20	24	0	20	100%	
-50% Cutoff	20	49	0	20	100%	
-25% Cutoff	20	74	1	19	95%	
+25%Cutoff	20	124	19	1	95%	
+50%Cutoff	20	148	20	0	100%	
+75%Cutoff	20	173	20	0	100%	

	Number of	Phencyclidine Concentration	Lay Perso	The Percentage	
% of Cutoff	Samples	by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	6	0	20	100%
-50% Cutoff	20	12.2	0	20	100%
-25% Cutoff	20	19	0	20	100%
+25%Cutoff	20	31.3	18	2	90%
+50%Cutoff	20	37	20	0	100%
+75%Cutoff	20	44	20	0	100%

	Number of	nples GC/MS (ng/mL)	Lay Person Results		The Percentage
% of Cutoff	Samples		No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	77	0	20	100%
-50% Cutoff	20	156	0	20	100%
-25% Cutoff	20	234	1	19	95%
+25%Cutoff	20	390	20	0	100%
+50%Cutoff	20	468	20	0	100%
+75%Cutoff	20	547	20	0	100%

	Number of	per of Buprenorphine Concentration	Lay Person Results		The Percentage
% of Cutoff	Samples	by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	2.6	0	20	100%
-50% Cutoff	20	5.2	0	20	100%
-25% Cutoff	20	7.8	1	19	95%
+25%Cutoff	20	13	19	1	95%
+50%Cutoff	20	15.7	20	0	100%
+75%Cutoff	20	18.3	20	0	100%

	Number of	of Morphine 300 Concentration Lay Pers	Lay Perso	on Results	The Percentage
% of Cutoff	Samples	by GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	74	0	20	100%
-50% Cutoff	20	148	0	20	100%
-25% Cutoff	20	228	0	20	100%
+25%Cutoff	20	379	19	1	95%
+50%Cutoff	20	443	20	0	100%
+75%Cutoff	20	516	20	0	100%

	Number of	EDDP Concentration by	Lay Person Results The P	The Percentage	
% of Cutoff	Samples GC/MS (ng/mL)	No. of Positive	No. of Negative	Agreement (%)	
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	81	0	20	100%
-50% Cutoff	20	157	0	20	100%
-25% Cutoff	20	235	1	19	95%
+25%Cutoff	20	410	20	0	100%
+50%Cutoff	20	485	20	0	100%
+75%Cutoff	20	566	20	0	100%

		Methylenedioxy- methamphetamine Concentration by GC/MS (ng/mL)	Lay Person Results		
% of Cutoff	Number of Samples		No. of Positive	No. of Positive	The Percentage Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	115	0	20	100%
-50% Cutoff	20	237	0	20	100%
-25% Cutoff	20	358	1	19	95%
+25%Cutoff	20	598	20	0	100%
+50%Cutoff	20	755	20	0	100%
+75%Cutoff	20	912	20	0	100%

	Number of	Nortriptyline Concentration by	Lay Person Results		The Percentage
% of Cutoff	Samples	GC/MS (ng/mL)	No. of Positive	No. of Positive	Agreement (%)
-100%Cutoff	20	0	0	20	100%
-75%Cutoff	20	261	0	20	100%
-50% Cutoff	20	495	0	20	100%
-25% Cutoff	20	720	1	19	95%
+25%Cutoff	20	1180	19	1	95%
+50%Cutoff	20	1485	20	0	100%
+75%Cutoff	20	1687	20	0	100%

#### BIBLIOGRAPHY OF SUGGESTED READING

- Stewart DJ, Inaba T, Lucassen M, Kalow W. Clin. Pharmacol. Ther. April 1979; 25 ed: 464, 264-8.
- 2. Ambre J. J. Anal. Toxicol. 1985; 9:241.
- 3. Hawks RL, CN Chiang. Urine Testing for Drugs of Abuse. National Institute for Drug Abuse (NIDA), Research Monograph 73, 1986.

ADDITIONAL INFORMATION AND REFERENCES

The following list of organizations may be helpful to you for counseling support and resources. These groups also have an Internet address, which can be access for additional information.

National Clearinghouse for Alcohol and Drug Information www.health.org 1-800-729-6686

Center for Substance Abuse Treatment www.health.org 1-800-662-HELP

The National Council on Alcoholism and Drug Dependence www.ncadd.org 1-800-NCA-CALL

American Council for Drug Education (ACDE) www.acde.org 1-800-488-DRUG

Manufactured for: W.H.P.M., Inc. 5358 Irwindale Ave. Irwindale, CA 91706 www.whpm.com

Effective Date: 12/12/2019